



1995 Updates:

Water Quality Criteria Documents for the Protection of Aquatic Life in Ambient Water

DISCLAIMER

This document has been reviewed by the Health and Ecological Criteria Division, Office of Science and Technology, U.S. Environmental Protection Agency, and approved for publication. Mention of trade names and commercial products does not constitute endorsement of their use.

ACKNOWLEDGMENT

Technical support for preparation of this document was provided to the Office of Water by Charles E. Stephan, U.S. EPA, Office of Research and Development, Mid-Continent Ecology Division, Duluth, MN.

CONTENTS

	<u>Page</u>
Arsenic (III)	A-1
Cadmium	B-1
Chromium (III)	C-1
Chromium (VI)	D-1
Copper	E-1
Cyanide	F-1
Dieldrin	G-1
Endrin	H-1
Lindane	I-1
Mercury (II)	J-1
Nickel	K-1
Parathion	L-1
Pentachlorophenol	M-1
Selenium	N-1
Zinc	O-1

INTRODUCTION

The purpose of these updates is to apply the methodology and datasets used in the derivation of the GLI aquatic life criteria to the national aquatic life criteria for these pollutants in fresh water. The methodology is that described for Tier I in Appendix A to Part 132: Great Lakes Water Quality Initiative Methodologies for Development of Aquatic Life Criteria and Values (Federal Register 60:15393-15399; March 23, 1995). This methodology differs from that described in the 1985 Guidelines (U.S. EPA 1985) in the following important ways:

- a. The GLI methodology gives preference to species that are resident in the Great Lakes System. This has no impact on these criteria, however, because the sensitive species in these datasets that are considered commercially or recreationally important for the purposes of deriving national aquatic life criteria are the same as the sensitive species in these datasets that are considered commercially or recreationally important for the purposes of deriving GLI aquatic life criteria.
- b. The GLI methodology does not use the Final Residue Value (FRV) that was used in the 1985 Guidelines. Instead of using the FRV in the derivation of aquatic life criteria, human health and wildlife criteria are to be derived using guidelines that are designed to provide adequate protection to human health and wildlife.
- c. Acute-Chronic Ratios (ACRs) for saltwater species are not used in the derivation of criteria for freshwater species if the Minimum Data Requirements for chronic data are satisfied by data for freshwater species.

Other aspects of the methodology are generally identical to those presented in the 1985 Guidelines.

Although it is not part of the methodology, if the range of Species Mean Acute Values (SMAVs) or Species Mean Chronic Values (SMCVs) within a genus was greater than a factor of five, the Genus Mean Acute Value or Genus Mean Chronic Value was set equal to the lowest SMAV or SMCV in that genus to provide adequate protection to the tested species in the genus. Whenever this was done, it is footnoted in the relevant table.

The datasets used in these updates used new data that were considered to be of acceptable quality along with the data in the criteria documents previously published by the U.S. EPA, which are referenced in the section for each pollutant. "New data" are data that became available since the last literature search used in the preparation of the criteria document by U.S. EPA and prior

to January 1993. Some errors in the U.S. EPA criteria documents were corrected and the new taxonomy for salmonids was used; some SMAVs and GMAVs are different from those in the U.S. EPA criteria documents due to the preference for results of "flow-through, measured" tests. Although some new data could have been used to revise the slopes relating acute and/or chronic toxicity to hardness or pH, it was decided that revision was not necessary at this time. Thus all of the slopes used herein are the same as those used in the criteria documents previously published by the U.S. EPA.

These updates affect criterion concentrations (i.e., Criterion Maximum Concentrations and/or Criterion Continuous Concentrations), but not averaging periods or frequencies of allowed exceedances. Four digits are given in the criterion concentrations because these are intermediate values in the derivation of permit limits.

The following abbreviations are used in this document:

ACR	= Acute-Chronic Ratio
CCC	= Criterion Continuous Concentration
CMC	= Criterion Maximum Concentration
FAV	= Final Acute Value
FCV	= Final Chronic Value
GMAV	= Genus Mean Acute Value
GMCV	= Genus Mean Chronic Value
FACR	= Final Acute-Chronic Ratio
SMACR	= Species Mean Acute-Chronic Ratio
SMAV	= Species Mean Acute Value
SMCV	= Species Mean Chronic Value

1995 UPDATE:
Freshwater Aquatic Life Criterion for Arsenic(III)

The new acceptable acute and chronic data for arsenic(III) are given in Tables A1 and A2. These new data were used with those given in Tables 1 and 2 of the criteria document for arsenic (U.S. EPA 1985) to obtain the values given in Table A3.

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values given in Table A3, resulting in a FAV of 679.6 ug/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 339.8 ug/L, as total recoverable arsenic(III).

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). The new chronic test gave an ACR of 3.784; the geometric mean of this value and the ACR in U.S. EPA (1985) for the same species was 4.199. This and the two other Species Mean ACRs in U.S. EPA (1985) are given in Table A3; the three ACRs were within a factor of 1.2. The FACR was calculated as the geometric mean of the three ACRs and was 4.594. The FCV = $FAV/FACR = (679.6 \text{ ug/L}) / (4.594) = 147.9 \text{ ug/L}$. This value did not need to be lowered to protect a commercially or recreationally important species. The CCC was 147.9 ug/L, as total recoverable arsenic(III).

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of arsenic(III) does not exceed 147.9 ug/L more than once every three years on the average and if the one-hour average concentration does not exceed 339.8 ug/L more than once every three years on the average.

Table A1. New Acute Values for Arsenic(III)

Species	Method*	Chemical	Test Duration (hrs)	Acute Value (ug/L)	Reference
Fathead minnow, <i>Pimephales promelas</i>	FT,M	Sodium arsenite	96	12,600	Spehar and Flandt 1986
Cladoceran, <i>Daphnia magna</i>	S,U	Sodium arsenite	48	4,501	Elnabarawy et al. 1986
Cladoceran, <i>Daphnia pulex</i>	S,U	Sodium arsenite	48	2,366	Elnabarawy et al. 1986
Cladoceran, <i>Ceriodaphnia reticulata</i>	S,U	Sodium arsenite	48	1,269	Elnabarawy et al. 1986

* FT = flow-through, M = measured, S = static, U = unmeasured.

Table A2. New Chronic Values for Arsenic(III)

Species	Test*	Acute Value (ug/L)	Chronic Value (ug/L)	Acute-Chronic Ratio	Reference
Fathead minnow, Pimephales promelas	ELS	12,600	3,330	3.784	Spehar and Fiantdt 1986

* ELS = early life stage.

Table A3. Ranked Genus Mean Acute Values for Arsenic(III)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
14	97,000	Midge, <i>Tanytarsus dissimilis</i>	97,000	-----
13	41,760	Bluegill, <i>Lepomis macrochirus</i>	41,760	-----
12	26,040	Goldfish, <i>Carassius auratus</i>	26,040	-----
11	24,500	Snail, <i>Aplexa hypnorum</i>	24,500	-----
10	22,040	Stonefly, <i>Pteronarcys californica</i>	22,040	-----
9	20,130	Flagfish, <i>Jordanella floridae</i>	20,130	4.862
8	18,100	Channel catfish <i>Ictalurus punctatus</i>	18,100	-----
7	14,960	Brook trout, <i>Salvelinus fontinalis</i>	14,960	-----
6	14,065	Fathead minnow, <i>Pimephales promelas</i>	14,065	4.199
5	13,340	Rainbow trout, <i>Oncorhynchus mykiss</i>	13,340	-----
4	2,690	Cladoceran, <i>Daphnia magna</i>	4,449	4.748
		Cladoceran, <i>Daphnia pulex</i>	1,626	-----
3	1,511	Cladoceran, <i>Ceriodaphnia reticulata</i>	1,511	-----
2	1,175	Cladoceran, <i>Simocephalus serrulatus</i>	812	-----
		Cladoceran, <i>Simocephalus vetulus</i>	1,700	-----
1	874	Amphipod, <i>Gammarus pseudolimnaeus</i>	874	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value. ∴

$$\text{FAV} = 679.6 \text{ ug/L}$$

$$\text{CMC} = \text{FAV}/2 = 339.8 \text{ ug/L}$$

$$\text{FACR} = 4.594$$

$$\text{FCV} = \text{FAV}/\text{FACR} = (679.6 \text{ ug/L})/(4.594) = 147.9 \text{ ug/L} = \text{CCC}$$

References

Elnabarawy, M.T., A.N. Welter, and R.R. Robideau. 1986. Relative Sensitivity of Three Daphnid Species to Selected Organic and Inorganic Chemicals. *Environ. Toxicol. Chem.* 5:393-398.

Spehar, R.L., and J.T. Fiandt. 1986. Acute and Chronic Effects of Water Quality Criteria-based Metal Mixtures on Three Aquatic Species. *Environ. Toxicol. Chem.* 5:917-931.

U.S. EPA. 1985. Ambient Water Quality Criteria for Arsenic - 1984. EPA 440/5-84-033. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Cadmium

The new acceptable acute and chronic data for cadmium are given in Tables B1 and B2. These new data were used with those given in Tables 1 and 2 of the criteria document for cadmium (U.S. EPA 1985) to obtain the values given in Tables B3 and B4. Because the toxicity of cadmium is hardness-dependent, all acute and chronic values in Tables B3 and B4 have been adjusted to a hardness of 50 mg/L.

Criterion Maximum Concentration (CMC)

The SMAVs given in Table B3 for the green sunfish, bluegill, coho salmon, and rainbow trout were derived from U.S. EPA (1985) by giving preference to results of "FT,M" tests. Several SMAVs given in U.S. EPA (1985) were changed or eliminated due to deletion of tests that were conducted in river water by Spehar and Carlson (1984a,b).

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values in Table B3, resulting in an FAV of 4.134 ug/L at a hardness of 50 mg/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 2.067 ug/L, as total recoverable cadmium, at a hardness of 50 mg/L. The CMC was related to hardness using the slope of 1.128 that was derived in U.S. EPA (1985):

$$\text{CMC} = e^{1.128 (\ln \text{hardness}) - 3.6867}$$

Criterion Continuous Concentration (CCC)

Two chronic values given in U.S. EPA (1985) were not used here because the tests were conducted in river water by Spehar and Carlson (1984a,b). The chronic value given in U.S. EPA (1985) for *Moina macrocopa* was not used here because the concentrations of cadmium were not measured.

Chronic toxicity tests have been conducted on cadmium with a wide variety of aquatic species and the resulting ACRs have a wide range, even within sensitive species (U.S. EPA 1985). Therefore, the Final Chronic Value (FCV) was calculated using the eight-family procedure that was used to calculate the FAV and was used to calculate the FCV for cadmium in U.S. EPA (1985). As in U.S.

EPA (1985), the FCV was calculated using the value of n used in the calculation of the FAV (i.e., n = 43). The FCV was 1.4286 ug/L at a hardness of 50 mg/L. This value did not need to be lowered to protect a commercially or recreationally important species. Thus the CCC was 1.4286 ug/L, as total recoverable cadmium, at a hardness of 50 mg/L. The CCC was related to hardness using the slope of 0.7852 that was derived in U.S. EPA (1985):

$$CCC = e^{0.7852 (\ln \text{ hardness}) - 2.715}$$

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of cadmium does not exceed the numerical value (in ug/L) given by the equation

$$CCC = e^{0.7852 (\ln \text{ hardness}) - 2.715}$$

more than once every three years on the average and if the one-hour average concentration does not exceed the numerical value (in ug/L) given by the equation

$$CMC = e^{1.128 (\ln \text{ hardness}) - 3.6867}$$

more than once every three years on the average.

Table B1. New Acute Values for Cadmium

Species	Method*	Hardness (mg/L as CaCO ₃)	Acute Value (ug/L)	Adjusted Acute Value (ug/L)**	Reference
Cladoceran, <i>Ceriodaphnia reticulata</i>	S,U	240	184	31.36	Elnabarawy et al. 1986
Cladoceran, <i>Daphnia pulex</i>	S,U	120	70	26.07	Hall et al. 1986
Cladoceran, <i>Daphnia pulex</i>	S,U	200	50	10.47	Hall et al. 1986
Cladoceran, <i>Daphnia pulex</i>	S,U	200	100	20.94	Hall et al. 1986
Cladoceran, <i>Daphnia pulex</i>	S,U	240	319	54.37	Elnabarawy et al. 1986
Cladoceran, <i>Daphnia magna</i>	S,U	240	178	30.3	Elnabarawy et al. 1986
Amphipod, <i>Crangonyx pseudogracilis</i>	S,U	50	1700	1700	Martin and Holdich 1986
Crayfish, <i>Orconectes virilis</i>	S,U	26	6100	12755	Mirenda 1986
Rainbow trout, <i>Oncorhynchus mykiss</i>	FT,M	9.2	<0.5	<3.37	Cusimano and Brakke 1986
Rainbow trout, <i>Oncorhynchus mykiss</i>	FT,M	50	30	30	Van Leeuwen et al. 1985
Rainbow trout, <i>Oncorhynchus mykiss</i>	FT,M	50	10	10	Van Leeuwen et al. 1985
Rainbow trout (28-day egg), <i>Oncorhynchus mykiss</i>	FT,M	50	9200	9200***	Van Leeuwen et al. 1985
Rainbow trout (14-day egg), <i>Oncorhynchus mykiss</i>	FT,M	50	7500	7500***	Van Leeuwen et al. 1985
Rainbow trout (24-hr. egg), <i>Oncorhynchus mykiss</i>	FT,M	50	13000	13000***	Van Leeuwen et al. 1985
Rainbow trout (0-hr. egg), <i>Oncorhynchus mykiss</i>	FT,M	50	13000	13000***	Van Leeuwen et al. 1985

Table B1. (Cont.)

Species	Method*	Hardness (mg/L as CaCO ₃)	Acute Value (ug/L)	Adjusted Acute Value (ug/L)**	Reference
Striped bass, Morone saxatilis	S,U	40	4	5.14	Palawski et al. 1985
Striped bass, Morone saxatilis	S,U	285	10	1.4	Palawski et al. 1985

* FT = flow-through, M = measured, S = static, U = unmeasured.

** Adjusted to a hardness of 50 mg/L using a slope of 1.128.

*** Not used in the calculation of the SMAV because data were available for a more sensitive life stage.

Table B2. New Chronic Values for Cadmium

Species	Test*	Hardness (mg/L as CaCO ₃)	Chronic Value (ug/L)	Adjusted Chronic Value (ug/L)**	Reference
Cladoceran, Ceriodaphnia reticulata	LC	240	0.4	0.12***	Elnabarawy et al. 1986
Cladoceran, Daphnia magna	LC	240	4.3	1.25***	Elnabarawy et al. 1986
Cladoceran, Daphnia pulex	LC	106	7.07	3.919	Ingersoll and Winner 1982
Cladoceran, Daphnia pulex	LC	65	7.49	6.096	Niederlehner 1984
Cladoceran, Daphnia pulex	LC	240	13.7	4***	Elnabarawy et al. 1986
Oligochaete, Aeolosoma headleyi	LC	65	25.19	20.50	Niederlehner 1984

* LC = life cycle.

** Adjusted to a hardness of 50 mg/L using a slope of 0.7852.

*** Not used in derivation of the criterion because the concentrations of cadmium were not measured.

Table B3. Ranked Genus Mean Acute Values for Cadmium

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**
43	12755	Crayfish, <i>Orconectes virilis</i>	12755
42	8325	Goldfish, <i>Carassius auratus</i>	8325
41	8100	Damselfly, (Unidentified)	8100
40	7921	Tubificid worm, <i>Rhyacodrilus montana</i>	7921
39	7685	Mosquitofish, <i>Gambusia affinis</i>	7685
38	6915	Tubificid worm, <i>Stylodrilus heringianus</i>	6915
37	4990	Tubificid worm, <i>Spirosperma ferox</i>	4401
		Tubificid worm, <i>Spirosperma nikolskyi</i>	5658
36	4977	Threespine stickleback <i>Gasterosteus aculeatus</i>	4977
35	4778	Tubificid worm, <i>Varichaeta pacifica</i>	4778
34	4024	Tubificid worm, <i>Tubifex tubifex</i>	4024
33	4024	Tubificid worm, <i>Quistradilus multisetosus</i>	4024
32	3800	Snail, <i>Amnicola</i> sp.	3800
31	3570	Guppy, <i>Poecilia reticulata</i>	3570
30	3514	White sucker, <i>Catostomus commersoni</i>	3514
29	3400	Caddisfly, (Unidentified)	3400
28	3018	Tubificid worm, <i>Branchiura sowerbyi</i>	3018

Table B3. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**
27	2888	Flagfish, <i>Jordanella floridae</i>	2888
26	2400	Northern squawfish, <i>Ptychocheilus oregonensis</i>	2400
25	2395	Green sunfish, <i>Lepomis cyanellus</i>	2399
		Pumpkinseed, <i>Lepomis gibbosus</i>	1347
		Bluegill, <i>Lepomis macrochirus</i>	4249
24	2310	Mayfly, <i>Ephemerella grandis</i>	2310
23	2137	Tubificid worm, <i>Limnodrilus hoffmeisteri</i>	2137
22	1700	Worm, <i>Nais</i> sp.	1700
21	1700	Amphipod, <i>Crangonyx pseudogracilis</i>	1700
20	1200	Midge, <i>Chironomus</i> sp.	1200
19	736	American eel, <i>Anguilla rostrata</i>	736
18	401	Isopod, <i>Asellus bicrenata</i>	401
17	221.9	Bryozoan, <i>Plumatella emarginata</i>	221.9
16	215.5	Common carp, <i>Cyprinus carpio</i>	215.5
15	156.9	Snail, <i>Physa gyrina</i>	156.9
14	142.5	Bryozoan, <i>Pectinatella magnifica</i>	142.5
13	104.0	Snail, <i>Aplexa hypnorum</i>	104.0

Table B3. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**
12	98.79	Banded killifish, <i>Fundulus diaphanus</i>	98.79
11	74.99	Amphipod, <i>Gammarus pseudolimnaeus</i>	80.33
		Amphipod, <i>Gammarus sp.</i>	70.00
10	48.28	Cladoceran, <i>Ceriodaphnia reticulata</i>	48.28
9	42.8	Isopod, <i>Lirceus alabamae</i>	42.8
8	40.78	Cladoceran, <i>Moina macrocopa</i>	40.78
7	30.54	Bryozoan, <i>Lophopodella carteri</i>	30.54
6	30.50	Fathead minnow, <i>Pimephales promelas</i>	30.50
5	29.96	Cladoceran, <i>Simocephalus serrulatus</i>	33.2
		Cladoceran, <i>Simocephalus vetulus</i>	27.03
4	21.13	Cladoceran, <i>Daphnia magna</i>	14.2
		Cladoceran, <i>Daphnia pulex</i>	31.43
3	5.421	Coho salmon, <i>Oncorhynchus kisutch</i>	6.48
		Chinook salmon, <i>Oncorhynchus tshawytscha</i>	4.254
		Rainbow trout, <i>Oncorhynchus mykiss</i>	5.78
2	2.682***	White perch, <i>Morone americana</i>	7544
		Striped bass, <i>Morone saxatilis</i>	2.682****

Table B3. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**
1	1.647	Brown trout, <i>Salmo trutta</i>	1.647

- * Ranked from most resistant to most sensitive based on Genus Mean Acute Value.
- ** At hardness = 50 mg/L.
- *** The GMAV was set equal to the lower SMAV due to the large range in the SMAVs in this genus.
- **** This SMAV was based on the results reported by Palawski et al. (1985) because they were considered better data than those given in U.S. EPA (1985), although the data reported by Hughes (1973) supported the newer data.

At hardness = 50 mg/L:

$$FAV = 4.134 \text{ ug/L}$$

$$CMC = FAV/2 = 2.067 \text{ ug/L}$$

As a function of hardness:

$$CMC = e^{1.128 (\ln \text{ hardness}) - 3.6867}$$

Table B4. Ranked Genus Mean Chronic Values for Cadmium

Rank*	Genus Mean Chronic Value (ug/L)**	Species	Species Mean Chronic Value (ug/L)**
12	20.50	Oligochaete, Aeolosoma headleyi	20.50
11	16.32	Bluegill, Lepomis macrochirus	16.32
10	15.40	Fathead minnow, Pimephales promelas	15.40
9	8.170	Smallmouth bass, Micropterus dolomieu	8.170
8	8.138	Northern pike, Esox lucius	8.138
7	7.849	White sucker, Catostomus commersoni	7.849
6	7.771	Atlantic salmon, Salmo salar	8.192
		Brown trout, Salmo trutta	7.372
5	5.336	Flagfish, Jordanella floridae	5.336
4	4.841	Snail, Aplexa hypnorum	4.841
3	4.383	Brook trout, Salvelinus fontinalis	2.362
		Lake trout, Salvelinus namaycush	8.134
2	3.399	Coho salmon, Oncorhynchus kisutch	4.289
		Chinook salmon, Oncorhynchus tshawytscha	2.694
1	0.1354***	Cladoceran, Daphnia magna	0.1354
		Cladoceran, Daphnia pulex	4.888

* Ranked from most resistant to most sensitive based on Genus Mean Chronic Value.

** At hardness = 50 mg/L.

*** The GMCV was set equal to the lower SMCV due to the large range in the SMCVs for this genus.

At hardness = 50 mg/L:

$$FCV = 1.4286 \text{ ug/L} = CCC \quad (\text{calculated using } n = 43)$$

As a function of hardness:

$$CCC = e^{0.7852 (\ln \text{ hardness}) - 2.715}$$

References

- Cusimano, R.F., D.F. Brakke, and G.A. Chapman. 1986. Effects of pH on the Toxicities of Cadmium, Copper and Zinc to Steelhead Trout (*Salmo gairdneri*). *Can. J. Fish. Aquat. Sci.* 43:1497-1503.
- Elnabarawy, M.R., A.N. Wetter, and R.R. Robideau. 1986. Relative Sensitivity of Three Daphnid Species to Selected Organic and Inorganic Chemicals. *Environ. Toxicol. Chem.* 5:393-398.
- Hall, W.S., R.L. Paulson, L.W. Hall, Jr., and D.T. Burton. 1986. Acute Toxicity of Cadmium and Sodium Pentachlorophenate to Daphnids and Fish. *Bull. Environ. Contam. Toxicol.* 37:308-316.
- Hughes, J.S. 1973. Acute Toxicity of Thirty Chemicals to Striped Bass (*Morone saxatilis*). Western Assoc. State Game Fish Comm., Salt Lake City, UT. July.
- Ingersoll, C.G., and R.W. Winner. 1982. Effect on *Daphnia pulex* (DeGeer) of Daily Pulse Exposures to Copper or Cadmium. *Environ. Toxicol. Chem.* 1:321-327.
- Martin, T.R., and D.M. Holdich. 1986. The Acute Lethal Toxicity of Heavy Metals to Peracarid Crustaceans (with Particular Reference to Fresh-water Asellids and Gammarids). *Water Res.* 20:1137-1147.
- Miranda, R.J. 1986. Toxicity and Accumulation of Cadmium in the Crayfish, *Orconectes virilis* (Hagen). *Arch. Environ. Contam. Toxicol.* 15:401-407.
- Niederlehner, B. 1984. A Comparison of Techniques for Estimating the Hazard of Chemicals in the Aquatic Environment. M.S. Thesis. Virginia Polytechnic Institute and State University.
- Palawski, D., J.B. Hunn, and F.J. Dwyer. 1985. Sensitivity of Young Striped Bass to Organic and Inorganic Contaminants in Fresh and Saline Waters. *Trans. Am. Fish. Soc.* 114:748-753.
- Spehar, R.L., and A.R. Carlson. 1984a. Derivation of Site-Specific Water Quality Criteria for Cadmium and the St. Louis River Basin, Duluth, Minnesota. PB84-153196. National Technical Information Service, Springfield, VA.
- Spehar, R.L., and A.R. Carlson. 1984b. Derivation of Site-Specific Water Quality Criteria for Cadmium and the St. Louis River Basin, Duluth, Minnesota. *Environ. Toxicol. Chem.* 3:651-665.

U.S. EPA. 1985. Ambient Aquatic Life Water Quality Criteria for Cadmium. EPA 440/5-84-032. National Technical Information Service, Springfield, VA.

Van Leeuwen, C.J., P.S. Griffioen, W.H.A. Vergouw, and J.L. Maas-Diepeveen. 1985. Differences in Susceptibility of Early Life Stages of Rainbow Trout (*Salmo gairdneri*) to Environmental Pollutants. *Aquatic Toxicol.* 7:59-78.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Chromium(III)

The new acceptable acute data for chromium(III) are given in Table C1; no new acceptable chronic data were found. These data were used with those given in Tables 1 and 2 of the criteria document for chromium (U.S. EPA 1984) to obtain the values given in Table C2. Because the toxicity of chromium(III) is hardness-dependent, all acute values in Table C2 have been adjusted to a hardness of 50 mg/L.

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values in Table C2, resulting in an FAV of 2044 ug/L at a hardness of 50 mg/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 1022 ug/L, as total recoverable chromium(III), at a hardness of 50 mg/L. The CMC was related to hardness using the slope of 0.819 that was derived in U.S. EPA (1985):

$$\text{CMC} = e^{0.819 (\ln \text{ hardness}) + 3.7256}$$

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). SMACRs were available for three species (Table C2) and the highest SMACR was obtained with the most resistant of the three. The other two SMACRs were within a factor of 2.4. The FACR was calculated as the geometric mean of the two ACRs and was 41.84. The FCV = FAV/FACR = (2044 ug/L)/(41.84) = 48.85 ug/L at a hardness of 50 mg/L. This value did not need to be lowered to protect a commercially or recreationally important species. Thus the CCC was 48.85 ug/L, as total recoverable chromium(III), at a hardness of 50 mg/L. The CCC, was related to hardness using the slope of 0.819:

$$\text{CCC} = e^{0.819 (\ln \text{ hardness}) + 0.6848}$$

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of chromium(III) does not exceed the numerical value (in ug/L) given by the equation

$$CCC = e^{0.819(\ln \text{ hardness}) + 0.6848}$$

more than once every three years on the average and if the one-hour average concentration does not exceed the numerical value (in ug/L) given by the equation

$$CMC = e^{0.819(\ln \text{ hardness}) + 3.7256}$$

more than once every three years on the average.

Table C1. New Acute Values for Chromium(III)

Species	Method*	Hardness (mg/L as CaCO ₃)	Acute Value (ug/L)	Adjusted Acute Value (ug/L)**	Reference
Amphipod, Crangonyx pseudogracilis	S,U	50	291,000	291,000	Martin and Holdich 1986

* S = static, U = unmeasured.

** Adjusted to a hardness of 50 mg/L using a slope of 0.819.

Table C2. Ranked Genus Mean Acute Values for Chromium(III)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
19	291,000	Amphipod, <i>Crangonyx pseudogracilis</i>	291,000	-----
18	71060	Caddisfly, <i>Hydropsyche betteni</i>	71060	-----
17	50000	Caddisfly, Unidentified sp.	50000	-----
16	43100	Damselfly, Unidentified sp.	43100	-----
15	16010	Cladoceran, <i>Daphnia magna</i>	16010	>356.4***
14	15630	Banded killifish, <i>Fundulus diaphanus</i>	15630	-----
13	15370	Pumpkinseed, <i>Lepomis gibbosus</i>	15720	-----
		Bluegill, <i>Lepomis macrochirus</i>	15020	-----
12	14770	White perch, <i>Morone americana</i>	13320	-----
		Striped bass, <i>Morone saxatilis</i>	16370	-----
11	13230	Common carp, <i>Cyprinus carpio</i>	13230	-----
10	12860	American eel, <i>Anguilla rostrata</i>	12860	-----
9	11000	Midge, <i>Chironomus sp.</i>	11000	-----
8	10320	Fathead minnow, <i>Pimephales promelas</i>	10320	27.30
7	10210	Snail, <i>Amnicola sp.</i>	10210	-----
6	9669	Rainbow trout, <i>Oncorhynchus mykiss</i>	9669	64.11
5	9300	Worm, <i>Nais sp.</i>	9300	-----

Table C2. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
4	8684	Goldfish, Carassius auratus	8684	-----
3	7053	Guppy, Poecilia reticulata	7053	-----
2	3200	Amphipod, Gammarus sp.	3200	-----
1	2221	Mayfly, Ephemerella subvaria	2221	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

** At hardness = 50 mg/L.

*** Not used in the calculation of the Final Acute-Chronic Ratio.

At hardness = 50 mg/L:

$$FAV = 2044 \text{ ug/L}$$

$$CMC = FAV/2 = 1022 \text{ ug/L}$$

As a function of hardness:

$$CMC = e^{0.819 (\ln \text{ hardness}) + 3.7256}$$

$$FACR = 41.84$$

At hardness = 50 mg/L:

$$FCV = FAV/FACR = (2044 \text{ mg/L}) / (41.84) = 48.85 \text{ ug/L} = CCC$$

As a function of hardness:

$$CCC = e^{0.819 (\ln \text{ hardness}) + 0.6848}$$

References

Martin, T.R., and D.M. Holdich. 1986. The Acute Lethal Toxicity of Heavy Metals to Peracarid Crustaceans (with Particular Reference to Fresh-water Asellids and Gammarids). Water Res. 20:1137-1147.

U.S. EPA. 1985. Ambient Aquatic Life Water Quality Criteria for Chromium(III) - 1984. EPA 440/5-84-029. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Chromium(VI)

The new acceptable acute data for chromium(VI) are given in Table D1; no new acceptable chronic data were used. These new data were used with those given in Tables 1 and 2 of the criteria document for chromium (U.S. EPA 1985) to obtain the values given in Table D2.

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values given in Table D2, resulting in a FAV of 32.04 ug/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 16.02 ug/L, as total recoverable chromium(VI).

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). Eight SMACRs were available (Table D2), but three were high SMACRs that were obtained with resistant species and one was a "greater than" value. Of the eight, only four were appropriate for use in calculating the FACR and the four were within a factor of 6. The FACR was calculated as the geometric mean of these four and was 2.917. The $FCV = FAV/FACR = (32.04 \text{ ug/L}) / (2.917) = 10.98 \text{ ug/L}$. This value did not need to be lowered to protect a commercially or recreationally important species. The CCC was 10.98 ug/L, as total recoverable chromium(VI).

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of chromium(VI) does not exceed 10.98 ug/L more than once every three years on the average and if the one-hour average concentration does not exceed 16.02 ug/L more than once every three years on the average.

Table D1. New Acute Values for Chromium(VI)

Species	Method*	Chemical	Acute Value (ug/L)	Reference
Cladoceran, Daphnia magna	S,U	K-dichromate	900**	Berglind and Dave 1984
Cladoceran, Daphnia magna	S,U	Na-dichromate	112**	Elnabarawy et al. 1986
Cladoceran, Daphnia pulex	S,M	K-dichromate	170**	Dorn et al. 1987
Cladoceran, Daphnia pulex	S,U	K-dichromate	190**	Dorn, et al. 1987
Cladoceran, Daphnia pulex	S,M	K-dichromate	20**	Dorn, et al. 1987
Cladoceran, Daphnia pulex	S,U	K-dichromate	20**	Dorn, et al. 1987
Cladoceran, Daphnia pulex	S,M	K-dichromate	40**	Dorn, et al. 1987
Cladoceran, Daphnia pulex	S,U	K-dichromate	40**	Dorn, et al. 1987
Cladoceran, Daphnia pulex	S,U	Na-dichromate	122**	Elnabarawy et al. 1986
Cladoceran, Daphnia pulex	S,M	K-dichromate	180**	Jop et al. 1987
Cladoceran, Daphnia pulex	S,M	K-dichromate	180**	Jop et al. 1987
Amphipod, Crangonyx pseudogracilis	R,U	K-dichromate	420	Martin and Holdich 1986
Amphipod, Crangonyx pseudogracilis	R,U	K-dichromate	810	Martin and Holdich 1986
Bluegill, Lepomis macrochirus	S,M	K-dichromate	182,000**	Jop et al. 1987
Bluegill, Lepomis macrochirus	S,M	K-dichromate	154,000**	Jop et al. 1987
Bluegill, Lepomis macrochirus	S,M	K-dichromate	201,240**	Dorn et al. 1987

Table D1. (Cont.)

Species	Method*	Chemical	Acute Value (ug/L)	Reference
Bluegill, Lepomis macrochirus	S,U	K-dichromate	164,730**	Dorn et al. 1987
Bluegill, Lepomis macrochirus	S,M	K-dichromate	199,200**	Dorn et al. 1987
Bluegill, Lepomis macrochirus	S,U	K-dichromate	158,360**	Dorn et al. 1987
Bluegill, Lepomis macrochirus	S,M	K-dichromate	148,310**	Dorn et al. 1987
Bluegill, Lepomis macrochirus	S,U	K-dichromate	146,530**	Dorn et al. 1987
Fathead minnow, Pimephales promelas	S,M	K-dichromate	46,000**	Jop et al. 1987
Fathead minnow, Pimephales promelas	S,M	K-dichromate	34,000**	Jop et al. 1987
Fathead minnow, Pimephales promelas	S,U	K-dichromate	26,130**	Dorn et al. 1987
Fathead minnow, Pimephales promelas	S,M	K-dichromate	26,410**	Dorn et al. 1987

* S = static, FT = flow-through, M = measured, U = unmeasured.

** Not used in the calculation of the SMAV because data were available for this species from a "FT,M" test.

Table D2. Ranked Genus Mean Acute Values for Chromium(VI)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
28	1,870,000	Stonefly, <i>Neophasganophora capitata</i>	1,870,000	-----
27	176,000	Crayfish, <i>Orconectes rusticus</i>	176,000	-----
26	140,000	Damselfly, <i>Enallagma aspersum</i>	140,000	-----
25	123,500	Green sunfish, <i>Lepomis cyanellus</i>	114,700	-----
		Bluegill, <i>Lepomis macrochirus</i>	132,900	-----
24	119,500	Goldfish, <i>Carassius auratus</i>	119,500	-----
23	72,600	White crappie, <i>Pomoxis annularis</i>	72,600	-----
22	69,000	Rainbow trout, <i>Oncorhynchus mykiss</i>	69,000	260.8**
21	67,610	Emerald shiner, <i>Notropis atherinoides</i>	48,400	-----
		Striped shiner, <i>Notropis chrysocephalus</i>	85,600	-----
		Sand shiner, <i>Notropis stramineus</i>	74,600	-----
20	61,000	Midge, <i>Chironomus tentans</i>	61,000	-----
19	59,000	Brook trout, <i>Salvelinus fontinalis</i>	59,000	223**
18	57,300	Midge, <i>Tanytarsus dissimilis</i>	57,300	-----
17	51,250	Central stoneroller, <i>Campostoma anomalum</i>	51,250	-----
16	49,600	Silverjaw minnow, <i>Ericymba buccata</i>	49,600	-----

Table D2. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
15	47,180	Bluntnose minnow, <i>Pimephales notatus</i>	54,225	-----
		Fathead minnow, <i>Pimephales promelas</i>	41,050	18.55**
14	46,000	Johnny darter, <i>Etheostoma nigrum</i>	46,000	-----
13	36,300	Yellow perch, <i>Perca flavescens</i>	36,300	-----
12	30,450	Striped bass, <i>Morone saxatilis</i>	30,450	-----
11	30,000	Guppy, <i>Poecilia reticulata</i>	30,000	-----
10	23,010	Snail, <i>Physa heterostropha</i>	23,010	-----
9	1,560	Bryozoan, <i>Lophopodella carteri</i>	1,560	-----
8	1,440	Bryozoan, <i>Pectinatella magnifica</i>	1,440	-----
7	650	Bryozoan, <i>Plumatella emarginata</i>	650	-----
6	630	Amphipod, <i>Hyalella azteca</i>	630	-----
5	583	Amphipod, <i>Crangonyx pseudogracilis</i>	583	-----
4	67.1	Amphipod, <i>Gammarus pseudolimnaeus</i>	67.1	-----
3	45.1	Cladoceran, <i>Ceriodaphnia reticulata</i>	45.1	1.13
2	36.35	Cladoceran, <i>Simocephalus serrulatus</i>	40.9	2.055
		Cladoceran, <i>Simocephalus vetulus</i>	32.3	5.267

Table D2. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
1	28.94	Cladoceran, Daphnia magna	23.07	>6.957**
		Cladoceran, Daphnia pulex	36.3	5.92

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

** Not used in the calculation of the Final Acute-Chronic Ratio.

$$\text{FAV} = 32.04 \text{ ug/L}$$

$$\text{CMC} = \text{FAV}/2 = 16.02 \text{ ug/L}$$

$$\text{FACR} = 2.917$$

$$\text{FCV} = \text{FAV}/\text{FACR} = (32.04 \text{ ug/L})/(2.917) = 10.98 \text{ ug/L} = \text{CCC}$$

References

Berglind, R., and G. Dave. 1984. Acute Toxicity of Chromate, DDT, PCP, TPBS, and Zinc to *Daphnia magna* Cultured in Hard and Soft Water. *Bull. Environ. Contam. Toxicol.* 33:63-68.

Dorn, P.B., J.H. Rodgers, Jr., K.M. Jop, J.C. Raia, and K.L. Dickson. 1987. Hexavalent Chromium as a Reference Toxicant in Effluent Toxicity Tests. *Environ. Toxicol. Chem.* 6:435-444.

Elnabarawy, M.T., A.N. Welter, and R.R. Robideau. 1986. Relative Sensitivity of Three Daphnid Species to Selected Organic and Inorganic Chemicals. *Environ. Toxicol. Chem.* 5:393-398.

Jop, K.M., T.F. Parkerton, J.H. Rodgers, and K.L. Dickson. 1987. Comparative Toxicity and Speciation of Two Hexavalent Chromium Salts in Acute Toxicity Tests. *Environ. Toxicol. Chem.* 6:697-703.

Martin, J.R., and D.M. Holdich. 1986. The Acute Lethal Toxicity of Heavy Metals to Peracarid Crustaceans (with Particular Reference to Fresh-water Asellids and Gammarids). *Water Res.* 20:1137-1147.

U.S. EPA. 1985. Ambient Water Quality Criteria for Chromium - 1984. EPA 440/5-84-029. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Copper

The new acceptable acute and chronic data for copper are given in Tables E1 and E2. These new data were used with those given in Tables 1 and 2 of the criteria document for copper (U.S. EPA 1985) to obtain the values given in Table E3. Because the toxicity of copper is hardness-dependent, all acute values in Table E3 have been adjusted to a hardness of 50 mg/L.

Criterion Maximum Concentration (CMC)

Data given in U.S. EPA (1985) for the species *Gammarus pulex* were not used because this species is not resident in North America. Several SMAVs given in Table E3 were derived from U.S. EPA (1985) by giving preference to results of "FT,M" tests.

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values in Table E3, resulting in an FAV of 14.57 ug/L at a hardness of 50 mg/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 7.285 ug/L, as total recoverable copper, at a hardness of 50 mg/L. The CMC was related to hardness using the slope of 0.9422 that was derived in U.S. EPA (1985):

$$\text{CMC} = e^{0.9422 (\ln \text{hardness}) - 1.700}$$

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). The new chronic test gave an ACR of 15.48 with the fathead minnow; the geometric mean of this value and the four ACRs for this species in U.S. EPA (1985) was 11.20. SMACRs were available for nine species (Table E3) and were higher for resistant species. To make the FACR appropriate for sensitive species, it was calculated from the two SMACRs that were determined with species whose SMAVs were close to the FAV. Thus the FACR was calculated as the geometric mean of 3.297 and 2.418 and was 2.823. The FCV = FAV/FACR = (14.57 ug/L)/(2.823) = 5.161 ug/L at a hardness of 50 mg/L. This value did not need to be lowered to protect a commercially or recreationally important species. Thus the CCC

was 5.161 ug/L, as total recoverable copper, at a hardness of 50 mg/L. The CCC was related to hardness using the slope of 0.8545 that was derived in U.S. EPA (1985):

$$CCC = e^{0.8545 (\ln \text{hardness}) - 1.702}$$

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of copper does not exceed the numerical value (in ug/L) given by the equation

$$CCC = e^{0.8545 (\ln \text{hardness}) - 1.702}$$

more than once every three years on the average and if the one-hour average concentration does not exceed the numerical value (in ug/L) given by the equation

$$CMC = e^{0.9422 (\ln \text{hardness}) - 1.700}$$

more than once every three years on the average.

Table E1. New Acute Values for Copper

Species	Method*	Hardness (mg/L as CaCO ₃)	Acute Value (ug/L)	Adjusted Acute Value (ug/L)**	Reference
Cladoceran, <i>Ceriodaphnia reticulata</i>	S,U	240	23	5.2	Elnabarawy et al. 1986
Cladoceran, <i>Daphnia magna</i>	S,U	240	41	9.4	Elnabarawy et al. 1986
Cladoceran, <i>Daphnia pulex</i>	S,U	240	31	7.1	Elnabarawy et al. 1986
Amphipod, <i>Crangonyx pseudogracilis</i>	S,U	50	1290	1290	Martin and Holdich 1986
Asiatic clam, <i>Corbicula manilensis</i>	FT,M	17	>2600	>7184	Harrison et al. 1984
Midge, <i>Chironomus decorus</i>	S,M	44	739	834	Kosalwat and Knight 1987
Fathead minnow, <i>Pimephales promelas</i>	FT,M	43.9	96	109	Spehar and Fiandt 1986
Bluegill, <i>Lepomis macrochirus</i>	S,M	31.2	340	530***	Bailey et al. 1985
Bluegill, <i>Lepomis macrochirus</i>	FT,M	31.2	550	858	Bailey et at. 1985
Rainbow trout, <i>Oncorhynchus mykiss</i>	FT,M	9.2	2.8	14	Cusimano and Brakke 1986
Striped bass, <i>Morone saxatilis</i>	S,U	285	270	52	Palawski et al. 1985

* S = static, FT = flow-through, U = unmeasured, M = measured.

** Adjusted to a hardness of 50 mg/L using the slope of 0.9422.

*** Not used in the calculation of the SMAV because data were available for this species from a "FT,M" test.

Table E2. New Chronic Values for Copper

Species	Test*	Acute Value (ug/L)	Chronic Value (ug/L)	Acute-Chronic Ratio	Reference
Fathead minnow, Pimephales promelas	ELS	96	6.2	15.48	Spehar and Fiandt 1986

* ELS = early life stage.

Table E3. Ranked Genus Mean Acute Values for Copper

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
43	10240	Stonefly, <i>Acroneuria lycorias</i>	10240	-----
42	> 7184	Asiatic clam, <i>Corbicula manilensis</i>	> 7184	-----
41	6200	Caddisfly, Unidentified sp.	6200	-----
40	4600	Damselfly, Unidentified sp.	4600	-----
39	4305	American eel, <i>Anguilla rostrata</i>	4305	-----
38	1990	Crayfish, <i>Procambarus clarkii</i>	1990	-----
37	1877	Snail, <i>Campeloma decisum</i>	1877	156.2***
36	1397	Crayfish, <i>Orconectes rusticus</i>	1397	-----
35	1290	Amphipod, <i>Crangonyx pseudogracilis</i>	1290	-----
34	1057	Pumpkinseed, <i>Lepomis gibbosus</i>	640.9	-----
		Bluegill, <i>Lepomis macrochirus</i>	1742	37.96***
33	900	Snail, <i>Amnicola</i> sp.	900	-----
32	790.6	Banded killifish, <i>Fundulus diaphanus</i>	790.6	-----
31	684.3	Mozambique tilapia <i>Tilapia mossambica</i>	684.3	-----
30	331.8	Striped shiner, <i>Notropis chrysocephalus</i>	331.8	-----
29	289	Goldfish, <i>Carassius auratus</i>	289	-----
28	242.7	Worm, <i>Lumbriculus variegatus</i>	242.7	-----

Table E3. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
27	196.1	Mosquitofish, <i>Gambusia affinis</i>	196.1	-----
26	170.2~	Midge, <i>Chironomus tentans</i>	197	-----
		Midge, <i>Chironomus decorus</i>	834	-----
		Midge, <i>Chironomus sp.</i>	30	-----
25	166.2	Snail, <i>Goniobasis livescens</i>	166.2	-----
24	156.8	Common carp, <i>Cyprinus carpio</i>	156.8	-----
23	141.2	Rainbow darter <i>Etheostoma caeruleum</i>	86.67	-----
		Orangethroat darter, <i>Etheostoma spectabile</i>	230.2	-----
22	135	Bryozoan, <i>Pectinatella magnifica</i>	135	-----
21	133	Chiselmouth, <i>Acrocheilus alutaceus</i>	133	-----
20	110.4	Brook trout, <i>Salvelinus fontinalis</i>	110.4	7.776***
19	109.9	Atlantic salmon, <i>Salmo salar</i>	109.9	-----
18	97.9	Bluntnose minnow, <i>Pimephales notatus</i>	72.16	26.36***
		Fathead minnow, <i>Pimephales promelas</i>	132.9	11.20***
17	90	Worm, <i>Nais sp.</i>	90	-----
16	86.67	Blacknose dace, <i>Rhinichthys atratulus</i>	86.67	-----

Table E3. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
15	83.97	Creek chub, <i>Semotilus atromaculatus</i>	83.97	-----
14	83	Guppy, <i>Poecilia reticulata</i>	83	-----
13	78.55	Central stoneroller, <i>Campostoma anomalum</i>	78.55	-----
12	73.99	Coho salmon, <i>Oncorhynchus kisutch</i>	87.1	-----
		Sockeye salmon, <i>Oncorhynchus nerka</i>	233.8	-----
		Cutthroat trout, <i>Oncorhynchus clarki</i>	66.26	-----
		Chinook salmon, <i>Oncorhynchus tshawytscha</i>	42.26	> 4.473***
		Rainbow trout, <i>Oncorhynchus mykiss</i>	38.89	-----
11	69.81	Brown bullhead, <i>Ictalurus nebulosus</i>	69.81	-----
10	56.21	Snail, <i>Gyraulus circumstriatus</i>	56.21	-----
9	53.08	Worm, <i>Limnodrilus hoffmeisteri</i>	53.08	-----
8	52~~	White perch, <i>Morone americanus</i>	5860	-----
		Striped bass, <i>Morone saxatilis</i>	52~~~	-----
7	39.33	Snail, <i>Physa heterostropha</i>	35.91	-----
		Snail, <i>Physa integra</i>	43.07	3.585***
6	37.05	Bryozoan, <i>Lophopodella carteri</i>	37.05	-----
5	37.05	Bryozoan, <i>Plumatella emarginata</i>	37.05	-----

Table E3. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
4	22.09	Amphipod, Gammarus pseudolimnaeus	22.09	3.297
3	16.74	Northern squawfish, Ptychocheilus oregonensis	16.74	-----
2	14.48	Cladoceran, Daphnia magna	19.88	2.418
		Cladoceran, Daphnia pulex	16.5	-----
		Cladoceran, Daphnia pulicaria	9.263	-----
1	9.92	Cladoceran, Ceriodaphnia reticulata	9.92	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

** At hardness = 50 mg/L.

*** Not used in the calculation of the Final Acute-Chronic Ratio.

~ This GMAV was not set equal to the lowest SMAV because the species was not identified and so might have been *C. tentans* or *C. decorus*.

~ This GMAV was set equal to the lower SMAV due to the large range in the SMAVs in this genus.

--- This SMAV was based on the results reported by Palawaki et al. (1985) because they were considered better data than those given in U.S. EPA (1985), although the data reported by Hughes (1973) supported the newer data.

At hardness = 50 mg/L:

$$FAV = 14.57 \text{ ug/L}$$

$$CMC = FAV/2 = 7.285 \text{ ug/L}$$

As a function of hardness:

$$CMC = e^{0.9422 (\ln \text{ hardness}) - 1.700}$$

$$FACR = 2.823$$

At hardness = 50 mg/L:

$$FCV = FAV/FACR = (14.57 \text{ ug/L}) / (2.823) = 5.161 \text{ ug/L} = CCC$$

As a function of hardness:

$$CCC = e^{0.8545 (\ln \text{ hardness}) - 1.702}$$

References

- Bailey, H.C., D.H.W. Liu, and H.A. Javitz. 1985. Time/Toxicity Relationships in Short-Term Static, Dynamic, and Plug-Flow Bioassays. In: Aquatic Toxicology and Hazard Assessment: Eighth Symposium. Bahner, R.C., and D.J. Hansen, Eds. ASTM STP 981. American Society for Testing and Materials, Philadelphia, PA. pp. 193-212.
- Cusimano, R.F., and D.F. Brakke. 1986. Effects of pH on the Toxicities of Cadmium, Copper, and Zinc to Steelhead Trout (*Salmo gairdneri*). Can. J. Fish. Aquat. Sci. 43:1497-1503.
- Elnabarawy, M.T., A.N. Welter, and R.R. Robideau. 1986. Relative Sensitivity of Three Daphnid Species to Selected Organic and Inorganic Chemicals. Environ. Toxicol. Chem. 5:393-398.
- Harrison, F.L., J.P. Knezovich, and D.W. Rice, Jr. 1984. The Toxicity of Copper to Adult and Early Life Stages of the Freshwater Clam, *Corbicula manilensis*. Arch. Environ. Contam. Toxicol. 13:85-92.
- Hughes, J.S. 1973. Acute Toxicity of Thirty Chemicals to Striped Bass (*Morone saxatilis*). Western Assoc. State Game Fish Comm., Salt Lake City, UT. July.
- Kosalwat, P., and A.W. Knight. 1987. Acute Toxicity of Aqueous and Substrate-Bound Copper to the Midge, *Chironomus decorus*. Arch. Environ. Contam. Toxicol. 16:275-282.
- Martin, T.R., and D.M. Holdich. 1986. The Acute Lethal Toxicity of Heavy Metals to Peracarid Crustaceans (with Particular Reference to Fresh-Water Asellids and Gammarids). Water Res. 20:1137-1147.
- Palawski, D., J.B. Hunn, and F.J. Dwyer. 1985. Sensitivity of Young Striped Bass to Organic and Inorganic Contaminants in Fresh and Saline Waters. Trans. Am. Fish. Soc. 114:748-753.
- Spehar, R.L., and J.T. Flandt. 1986. Acute and Chronic Effects of Water Quality Criteria-Based Metal Mixtures on Three Aquatic Species. Environ. Toxicol. Chem. 5:917-931.
- U.S. EPA. 1985. Ambient Aquatic Life Water Quality Criteria for Copper. EPA 440/5-84-031. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Cyanide

No new acceptable acute or chronic data were found for cyanide. Therefore, the data in the existing criteria document for cyanide (U.S. EPA 1985) were used as the basis for the derivation of this criterion. The new taxonomy for salmonids was used (Table F1), but this did not cause a change in the criterion for cyanide.

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values given in Table F1, resulting in a FAV of 45.77 ug/L. Because the SMAV of the commercially and recreationally important rainbow trout was 44.73 ug/L, the FAV was lowered to 44.73 ug/L. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 22.36 ug free cyanide (as CN)/L.

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). Five SMACRs are available (Table F1), but one was a high SMACR that was obtained with a resistant species; the other four were within a factor of 1.5. The FACR was calculated as the geometric mean of these four and was 8.568. The $FCV = FAV/FACR = (44.73 \text{ ug/L}) / (8.568) = 5.221 \text{ ug/L}$. This value does not need to be lowered to protect a commercially or recreationally important species. The CCC was 5.221 ug free cyanide (as CN)/L.

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of free cyanide (as CN) does not exceed 5.221 ug/L more than once every three years on the average and if the one-hour average concentration does not exceed 22.36 ug/L more than once every three years on the average.

Table F1. Ranked Genus Mean Acute Values for Cyanide

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
16	2490	Midge, <i>Tanytarsus dissimilis</i>	2490	-----
15	2326	Isopod, <i>Asellus communis</i>	2326	68.29**
14	432	Snail, <i>Physa heterostropha</i>	432	-----
13	426	Stonefly, <i>Pteronarcys dorsata</i>	426	-----
12	318	Goldfish, <i>Carassius auratus</i>	318	-----
11	167	Amphipod, <i>Gammarus pseudolimnaeus</i>	167	9.111
10	147	Guppy, <i>Poecilia reticulata</i>	147	-----
9	125.1	Fathead minnow, <i>Pimephales promelas</i>	125.1	7.633
8	123.6	Cladoceran, <i>Daphnia magna</i>	160	-----
		Cladoceran, <i>Daphnia pulex</i>	95.55	-----
7	102	Largemouth bass, <i>Micropterus salmoides</i>	102	-----
6	102	Black crappie, <i>Pomoxis nigromaculatus</i>	102	-----
5	99.28	Bluegill, <i>Lepomis macrochirus</i>	99.28	7.316
4	92.64	Yellow perch, <i>Perca flavescens</i>	92.64	-----
3	90.00	Atlantic salmon, <i>Salmo salar</i>	90.00	-----
2	85.80	Brook trout, <i>Salvelinus fontinalis</i>	85.80	10.59
1	44.73	Rainbow trout <i>Oncorhynchus mykiss</i>	44.73	-----

-
- * Ranked from most resistant to most sensitive based on Genus Mean Acute Value.
 - ** Not used in the calculation of the Final Acute-Chronic Ratio.

Calculated FAV = 45.77 ug/L

Lowered to protect rainbow trout:

$$\text{FAV} = 44.73 \text{ ug/L}$$

$$\text{CMC} = \text{FAV}/2 = 22.36 \text{ ug/L}$$

$$\text{FACR} = 8.568$$

$$\text{FCV} = \text{FAV}/\text{FACR} = (44.73 \text{ ug/L}) / (8.568) = 5.221 \text{ ug/L} = \text{CCC}$$

References

U.S. EPA. 1985. Ambient Water Quality Criteria for Cyanide - 1984. EPA 440/5-84-028. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Dieldrin

The new acceptable acute data for dieldrin are given in Table G1; no new acceptable chronic data were found. These new data were used with those given in Tables 1 and 2 of the criteria document for dieldrin (U.S. EPA 1980) to obtain the values given in Table G2. Although results from the following publications were used in U.S. EPA (1980), they were not considered acceptable for use here: Santharam et al. (1976), Gaufin (1965), and Jensen and Gaufin (1964).

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values given in Table G2, resulting in a FAV of 0.4749 ug/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 0.2374 ug/L.

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). Two SMACRs were given in Table G2; a third SMACR of 6.2 was given in U.S. EPA (1980) for the saltwater mysid. These three were within a factor of 1.8. The FACR was calculated as the geometric mean of the three SMACRs and was 8.530. The FCV = $FAV/FACR = (0.4749 \text{ ug/L})/(8.530) = 0.0557 \text{ ug/L}$. This value did not need to be lowered to protect a commercially or recreationally important species. The CCC was 0.0557 ug/L.

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of dieldrin does not exceed 0.0557 ug/L more than once every three years on the average and if the one-hour average concentration does not exceed 0.2374 ug/L more than once every three years on the average.

Table G1. New Acute Values for Dieldrin

Species	Method*	Test Duration (hrs)	Acute Value (ug/L)	Reference
Cladoceran, <i>Daphnia pulex</i>	S,M	48	251	Daniels and Allan 1981
Cladoceran, <i>Daphnia pulex</i>	S,U	48	190	Mayer and Ellersieck 1986
Stonefly, <i>Claassenia sabulosa</i>	S,U	96	0.6	Mayer and Ellersieck 1986
Stonefly, <i>Pteronarcys californica</i>	S,U	96	0.5	Mayer and Ellersieck 1986
Stonefly, <i>Pteronarcella badia</i>	S,U	96	0.5	Mayer and Ellersieck 1986
Damselfly, <i>Ischnura verticalis</i>	S,U	96	12	Mayer and Ellersieck 1986
Annelid, <i>Lumbriculus variegatus</i>	FT,M	96	21.8	Brooke 1993
Rainbow trout, <i>Oncorhynchus mykiss</i>	S,U	96	1.2**	Mayer and Ellersieck 1986
Rainbow trout, <i>Oncorhynchus mykiss</i>	FT,M	96	0.62	Shubat and Curtis 1986
Rainbow trout, <i>Oncorhynchus mykiss</i>	S,U	96	3**	Van Leeuwen et al. 1985
Goldfish, <i>Carassius auratus</i>	S,U	96	1.8	Mayer and Ellersieck 1986
Fathead minnow, <i>Pimephales promelas</i>	S,U	96	3.8	Mayer and Ellersieck 1986
Bluegill, <i>Lepomis macrochirus</i>	S,U	96	3.1	Mayer and Ellersieck 1986
Bluegill, <i>Lepomis macrochirus</i>	S,U	96	7	Sanders 1972
Pumpkinseed, <i>Lepomis gibbosus</i>	S,U	96	6.7	Cairns and Scheier 1964
Cutthroat trout, <i>Oncorhynchus clarki</i>	S,U	96	6	Mayer and Ellersieck 1986

Table G1. (Cont.)

Species	Method*	Test Duration (hrs)	Acute Value (ug/L)	Reference
Channel catfish, <i>Ictalurus punctatus</i>	S,U	96	4.5	Mayer and Ellersieck 1986
Largemouth bass, <i>Micropterus salmoides</i>	S,U	96	3.5	Mayer and Ellersieck 1986

* S = static, FT = flow-through, U = unmeasured, M = measured.

** Not used in the calculation of the SMAV because data were available for this species from a "FT,M" test.

Table G2. Ranked Genus Mean Acute Values for Dieldrin

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
18	740	Crayfish, Orconectes nais	740	-----
17	534	Amphipod, Gammarus lacustris	460	-----
		Amphipod, Gammarus fasciatus	620	-----
16	228	Cladoceran, Daphnia pulex	228	-----
15	214	Cladoceran, Simocephalus serrulatus	214	-----
14	21.8	Annelid, Lumbriculus variegatus	21.8	-----
13	20	Glass shrimp Palaemonetes kadiakensis	20	-----
12	17.7	Fathead minnow, Pimephales promelas	17.7	-----
11	12	Damselfly, Ischnura verticalis	12	-----
10	8.6	Goldfish, Carassius auratus	8.6	-----
9	8.5	Pumpkinseed, Lepomis gibbosus	6.7	-----
		Bluegill, Lepomis macrochirus	11.5	-----
		Green sunfish, Lepomis cyanellus	8.1	-----
8	5	Isopod, Asellus brevicaudus	5	-----
7	4.5	Channel catfish, Ictalurus punctatus	4.5	-----
6	4.5	Guppy, Poecilia reticulata	4.5	9.1

Table G2. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
5	3.5	Largemouth bass, Micropterus salmoides	3.5	-----
4	0.62**	Chinook salmon, Oncorhynchus tshawytscha	6.1	-----
		Coho salmon, Oncorhynchus kisutch	10.8	-----
		Cutthroat trout, Oncorhynchus clarki	6	-----
		Rainbow trout, Oncorhynchus mykiss	0.62	11
3	0.6	Stonefly, Claassenia sabulosa	0.6	-----
2	0.5	Stonefly, Pteronarcys californica	0.5	-----
1	0.5	Stonefly, Pteronarcella badia	0.5	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

** The GMAV was set equal to the lowest SMAV due to the large range in the SMAVs in this genus.

$$FAV = 0.4749 \text{ ug/L}$$

$$CMC = FAV/2 = 0.2374 \text{ ug/L}$$

$$FACR = 8.530$$

$$FCV = FAV/FACR = (0.4749 \text{ ug/L}) / (8.530) = 0.0557 \text{ ug/L} = CCC$$

References

Brooke, L. 1993. Acute and Chronic Toxicity of Several Pesticides to Five Species of Aquatic Organisms. Report to R. Spehar, U.S. EPA, Duluth, MN. 31 pp.

Cairns, J., Jr., and A. Scheier. 1964. The Effect on the Sunfish, *Lepomis gibbosus*, of Chronic Exposure to Lethal and Sublethal Concentrations of Dieldrin. *Notulae Naturae* 310:1-10.

Daniels, R.E., and J.D. Allan. 1981. Life Table Evaluation of Chronic Exposure to a Pesticide. *Can. J. Fish. Aquat. Sci.* 38:485-494.

Mayer, F.L., and M.R. Ellersieck. 1986. Manual of Acute Toxicity: Interpretation and Data Base for 410 Chemicals and 66 Species of Fresh-water Animals. USDI Publication 160. Columbia, MO.

Sanders, H.O. 1972. Toxicity of Some Insecticides to Four Species of Malacostracan Crustaceans. U.S. Bureau Sports Fishery and Wildlife Tech. Paper 66.

Shubat, P.J., and L.R. Curtis. 1986. Ration and Toxicant Preexposure Influence on Dieldrin Accumulation by Rainbow Trout (*Salmo gairdneri*). *Environ. Toxicol. Chem.* 5:69-77.

U.S. EPA. 1980. Ambient Water Quality Criteria for Aldrin/Dieldrin. EPA-440/5-80-019. National Technical Information Service, Springfield, VA.

Van Leeuwen, C.J., P.S. Griffioen, W.H.A. Vergouw, and J.L. Maas-Diepeveen. 1985. Differences in Susceptibility of Early Life Stages of Rainbow Trout (*Salmo gairdneri*) to Environmental Pollutants. *Aquatic Toxicol.* 7:59-78.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Endrin

The new acceptable acute data for endrin are given in Table H1; no new acceptable chronic data were found. These new data were used with those given in Tables 1 and 2 of the criteria document for endrin (U.S. EPA 1980) to obtain the values given in Table H2. Results in the following publications were used in U.S. EPA (1980) but were not considered acceptable for use here: Katz and Chadwick (1961), Naqui and Ferguson (1968), Nebeker and Gaufin (1964), Gaufin et al. (1965), Jensen and Gaufin (1966), Post and Schroeder (1971), Mount (1962), and Solon (1969).

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values given in Table H2, resulting in a FAV of 0.1728 ug/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 0.0864 ug/L.

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). Four ACRs were given in U.S. EPA (1980) but the ACR for the fathead minnow was considered unacceptable for use here. ACRs of 1.9 and 18 were determined with saltwater species, whereas an ACR of 3.3 was obtained with a freshwater species (Table H2); the three were within a factor of 9.5. The FACR was calculated as the geometric mean of the other three and was 4.833. The FCV = FAV/FACR = $(0.1728 \text{ ug/L}) / (4.833) = 0.03575 \text{ ug/L}$. This value did not need to be lowered to protect a commercially or recreationally important species. The CCC was 0.03575 ug/L.

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of endrin does not exceed 0.03575 ug/L more than once every three years on the average and

if the one-hour average concentration does not exceed 0.0864 ug/L more than once every three years on the average.

Table H1. New Acute Values for Endrin

Species	Method*	Test Duration (hrs)	Acute Value (ug/L)	Reference
Cladoceran, <i>Ceriodaphnia reticulata</i>	S,U	48	24	Elnabarawy et al. 1986
Cladoceran, <i>Daphnia magna</i>	S,U	48	4.2	Mayer and Ellersieck 1985
Cladoceran, <i>Daphnia magna</i>	S,U	48	59	Elnabarawy et al. 1986
Cladoceran, <i>Daphnia magna</i>	S,U	48	41	Mayer and Ellersieck 1985
Cladoceran, <i>Daphnia magna</i>	S,U	48	74	Mayer and Ellersieck 1985
Cladoceran, <i>Daphnia magna</i>	S,M	48	160	Thurston et al. 1985
Cladoceran, <i>Daphnia pulex</i>	S,U	48	20	Mayer and Ellersieck 1985
Cladoceran, <i>Daphnia pulex</i>	S,U	48	30	Elnabarawy et al. 1986
Annelid, <i>Lumbriculus variegatus</i>	FT,M	36	42.6	Brooke 1993
Snipe fly, <i>Atherix variegatus</i>	S,U	96	4.6	Mayer and Ellersieck 1985
Midge, <i>Tanytarsus dissimilis</i>	S,M	48	0.84	Thurston et al. 1985
Stonefly, <i>Acronuria pacifica</i>	S,U	96	> 0.18**	Mayer and Ellersieck 1985
Crayfish, <i>Orconectes immunis</i>	FT,M	96	89	Thurston et al. 1985
Damselfly, <i>Ischnura verticalis</i>	S,U	96	2.4	Mayer and Ellersieck 1986
Damselfly, <i>Ischnura verticalis</i>	S,U	96	2.1	Mayer and Ellersieck 1986
Yellow perch, <i>Perca flavescens</i>	FT,U	96	0.15	Mayer and Ellersieck 1986
Largemouth bass, <i>Micropterus salmoides</i>	S,U	96	0.31	Mayer and Ellersieck 1986

Table H1. (Cont.)

Species	Method*	Test Duration (hrs)	Acute Value (ug/L)	Reference
Black bullhead, <i>Ictalurus melas</i>	S,U	96	1.1	Mayer and Ellersieck 1986
Channel catfish, <i>Ictalurus punctatus</i>	S,U	96	0.32***	Mayer and Ellersieck 1986
Channel catfish, <i>Ictalurus punctatus</i>	S,U	96	1.1***	Mayer and Ellersieck 1986
Channel catfish, <i>Ictalurus punctatus</i>	FT,M	96	0.42	Thurston et al. 1985
Rainbow trout, <i>Oncorhynchus mykiss</i>	S,U	96	0.75***	Mayer and Ellersieck 1986
Rainbow trout, <i>Oncorhynchus mykiss</i>	FT,M	96	0.3	Thurston et al. 1985
Goldfish, <i>Carassius auratus</i>	FT,U	96	0.44***	Mayer and Ellersieck 1986
Goldfish, <i>Carassius auratus</i>	FT,M	96	0.95	Thurston et al. 1985
Fathead minnow, <i>Pimephales promelas</i>	S,U	96	1.8***	Mayer and Ellersieck 1986
Fathead minnow, <i>Pimephales promelas</i>	FT,M	96	0.65	Thurston et al. 1985
Mosquitofish, <i>Gambusia affinis</i>	S,U	96	1.1***	Mayer and Ellersieck 1986
Mosquitofish, <i>Gambusia affinis</i>	FT,M	96	0.69	Thurston et al. 1985
Carp, <i>Cyprinus carpio</i>	FT,U	96	0.32	Mayer and Ellersieck 1986
Bluegill, <i>Lepomis macrochirus</i>	FT,M	96	0.21	Thurston et al. 1985
Bullfrog tadpole, <i>Rana catesbeiana</i>	FT,M	96	2.5	Thurston et al. 1985

* FT = flow-through, S = static, U = unmeasured, M = measured.

** Not used in the calculation of the FAV because it is not appropriate to have one of the four lowest GMAVs be a "greater than" value.

*** Not used in the calculation of the SMAV because data were available for this species from a "FT,M" test.

Table H2. Ranked Genus Mean Acute Values for Endrin

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
27	64	Mayfly, <i>Hexagenia bilineata</i>	64	-----
26	53	Crayfish, <i>Orconectes nais</i>	32	-----
		Crayfish, <i>Orconectes immunis</i>	89	-----
25	43	Annelid, <i>Lumbriculus variegatus</i>	43	-----
24	38	Cladoceran, <i>Daphnia magna</i>	59	-----
		Cladoceran, <i>Daphnia pulex</i>	24	-----
23	34	Cladoceran, <i>Simocephalus serrulatus</i>	34	-----
22	24	Cladoceran, <i>Ceriodaphnia reticulata</i>	24	-----
21	4.6	Snipe fly, <i>Atherix variegatus</i>	4.6	-----
20	3.0	Amphipod, <i>Gammarus fasciatus</i>	3.1	-----
		Amphipod, <i>Gammarus lacustris</i>	3.0	-----
19	2.5	Bullfrog tadpole <i>Rana catesbeiana</i>	2.5	-----
18	2.1	Damselfly, <i>Ischnura verticalis</i>	2.1	-----
17	1.6	Guppy, <i>Poecilia reticulata</i>	1.6	-----
16	1.5	Isopod, <i>Asellus brevicaudus</i>	1.5	-----
15	1.3	Glass shrimp, <i>Palaemonetes kadiakensis</i>	1.3	-----
14	0.95	Goldfish, <i>Carassius auratus</i>	0.95	-----

Table H2. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
13	0.85	Flagfish, <i>Jordanella floridae</i>	0.85	3.3
12	0.84	Midge, <i>Tanytarsus dissimilis</i>	0.84	-----
11	0.76	Stonefly, <i>Claassenia sabulosa</i>	0.76	-----
10	0.69	Mosquitofish, <i>Gambusia affinis</i>	0.69	-----
9	0.68	Black bullhead, <i>Ictalurus melas</i>	1.1	-----
		Channel catfish, <i>Ictalurus punctatus</i>	0.42	-----
8	0.57	Coho salmon, <i>Oncorhynchus kisutch</i>	0.51	-----
		Chinook salmon, <i>Oncorhynchus tshawytscha</i>	1.2	-----
		Rainbow trout, <i>Oncorhynchus mykiss</i>	0.3	-----
7	0.54	Stonefly, <i>Pteronarcella badia</i>	0.54	-----
6	0.49	Fathead minnow, <i>Pimephales promelas</i>	0.49	-----
5	0.32	Common carp, <i>Cyprinus carpio</i>	0.32	-----
4	0.31	Largemouth bass, <i>Micropterus salmoides</i>	0.31	-----
3	0.25	Stonefly, <i>Pteronarcys californica</i>	0.25	-----
2	0.21	Bluegill, <i>Lepomis macrochirus</i>	0.21	-----
1	0.15	Yellow perch, <i>Perca flavescens</i>	0.15	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

$$\text{FAV} = 0.1728 \text{ ug/L}$$

$$\text{CMC} = \text{FAV}/2 = 0.0864 \text{ ug/L}$$

$$\text{FACR} = 4.833$$

$$\text{FCV} = \text{FAV}/\text{FACR} = (0.1728 \text{ ug/L})/(4.833) = 0.03575 \text{ ug/L} = \text{CCC}$$

References

Brooke, L. 1993. Acute and Chronic Toxicity of Several Pesticides to Five Species of Aquatic Organisms. Report to R. Spehar, U.S. EPA, Duluth, MN. 31 pp.

Elnabarawy, M.T., A.N. Welter, and R.R. Robideau. 1986. Relative Sensitivity of Three Daphnid Species to Selected Organic and Inorganic Chemicals. Environ. Toxicol. Chem. 5:393-398.

Mayer, F.L., and M.R. Ellersieck. 1986. Manual of Acute Toxicity: Interpretation and Data Base for 410 Chemicals and 66 Species of Freshwater Animals. USDI Publication 160. Columbia, MO.

Thurston, R.V., T.A. Gilfoil, E.L. Meyn, R.K. Zajdel, T.I. Aoki and G.D. Veith. 1985. Comparative Toxicity of Ten Organic Chemicals to Ten Common Aquatic Species. Water Res. 19:1145-1155.

U.S. EPA. 1980. Ambient Water Quality Criteria for Endrin. EPA 440/5-80-047. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Lindane

The new acceptable acute data for lindane are given in Table I1; no new acceptable chronic data were found. These new data were used with those given in Tables 1 and 2 of the criteria document for lindane (U.S. EPA 1980) to obtain the values given in Table I2.

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values given in Table I2, resulting in a FAV of 1.903 ug/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 0.9515 ug/L.

Criterion Continuous Concentration (CCC)

Three ACRs were given in U.S. EPA (1980) but the ACR for the fathead minnow was considered unacceptable for use here. No new ACRs were available and so a FCV could not be calculated using either the eight-family procedure or the FACR procedure. Therefore, a CCC could not be determined.

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably by acute toxicity if the one-hour average concentration of lindane does not exceed 0.9515 ug/L more than once every three years on the average.

Table II. New Acute Values for Lindane

Species	Method*	Test Duration (hrs)	Acute Value (ug/L)	Reference
Cladoceran, Daphnia magna	S,U	48	516	Randall et al. 1979
Cladoceran, Daphnia magna	S,M	48	1000	Hermens et al. 1984
Amphipod, Gammarus lacustris	S,U	96	88	Mayer and Ellersieck 1986
Snail, Lymnaea stagnalis	S,U	96	3.3	Bluzat and Senge 1979
Stonefly, Pteronarcys californicus	S,U	96	4.5	Mayer and Ellersieck 1986
Stonefly, Pteronarcys californicus	S,U	96	1	Mayer and Ellersieck 1986
Damselfly, Lestes congener	S,U	96	20	Federle and Collins 1976
Backswimmer, Notonecta undulata	S,U	96	3	Federle and Collins 1976
Crawling water beetle, Peltodytes sp.	S,U	96	20	Federle and Collins 1976
Coho salmon, Oncorhynchus kisutch	S,U	96	23	Mayer and Ellersieck 1986
Lake trout, Salvelinus namaycush	S,U	96	32	Mayer and Ellersieck 1986
Lake trout, Salvelinus namaycush	S,U	96	24	Mayer and Ellersieck 1986
Brown trout, Salmo trutta	S,U	96	24	Mayer and Ellersieck 1986
Brown trout, Salmo trutta	S,U	96	25	Mayer and Ellersieck 1986
Brown trout, Salmo trutta	FT,U	96	22	Mayer and Ellersieck 1986
Rainbow trout, Oncorhynchus mykiss	FT,M	96	22	Tooby and Durbin 1975
Rainbow trout, Oncorhynchus mykiss	S,U	96	18**	Mayer and Ellersieck 1986

Table II. (Cont).

Species	Method*	Test Duration (hrs)	Acute Value (ug/L)	Reference
Rainbow trout, <i>Oncorhynchus mykiss</i>	S,U	96	24**	Mayer and Ellersieck 1986
Rainbow trout, <i>Oncorhynchus mykiss</i>	S,U	96	31**	Mayer and Ellersieck 1986
Rainbow trout, <i>Oncorhynchus mykiss</i>	S,U	96	41**	Mayer and Ellersieck 1986
Rainbow trout, <i>Oncorhynchus mykiss</i>	FT,M	96	30	Tooby and Durbin 1975
Bluegill, <i>Lepomis macrochirus</i>	S,U	96	57	Randall et al. 1979
Bluegill, <i>Lepomis macrochirus</i>	S,U	96	56	Mayer and Ellersieck 1986
Green sunfish, <i>Lepomis cyanellus</i>	S,U	96	70	Mayer and Ellersieck, 1986
Green sunfish, <i>Lepomis cyanellus</i>	S,U	96	83	Mayer and Ellersieck 1986
Yellow perch, <i>Perca flavescens</i>	FT,U	96	23	Mayer and Ellersieck 1986
Fathead minnow, <i>Pimephales promelas</i>	FT,U	96	77	Mayer and Ellersieck 1986
Fathead minnow, <i>Pimephales promelas</i>	S,U	96	67	Mayer and Ellersieck 1986
Fathead minnow, <i>Pimephales promelas</i>	S,U	96	86	Mayer and Ellersieck 1986
Goldfish, <i>Carassius auratus</i>	S,U	96	90	Macek and McAllister 1970
Goldfish, <i>Carassius auratus</i>	S,U	96	105	Mayer and Ellersieck 1986
Channel catfish, <i>Ictalurus punctatus</i>	S,U	96	49	Mayer and Ellersieck 1986
Fowlers toad, <i>Bufo woodhousei fowleri</i>	S,U	96	3200	Mayer and Ellersieck 1986
Western chorus frog, <i>Pseudacris triseriata</i>	S,U	96	2650	Mayer and Ellersieck 1986

-
- * S = static, FT = flow-through, U = unmeasured, M = measured.
 - ** Not used in the calculation of the SMAV because data were available for this species from a "FT,M" test.

Table I2. Ranked Genus Mean Acute Values for Lindane

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
23	3200	Fowlers toad, Bufo woodhousi fowleri	3200	-----
22	2650	Western chorus frog, Pseudacris triseriata	2650	-----
21	676	Cladoceran, Simocephalus serrulatus	676	-----
20	538	Cladoceran, Daphnia magna	630	33
		Cladoceran, Daphnia pulex	460	-----
19	207	Midge, Chironomus tentans	207	63
18	138	Guppy, Poecilia reticulata	138	-----
17	117	Goldfish, Carassius auratus	117	-----
16	90	Carp, Cyprinus carpio	90	-----
15	72	Fathead minnow, Pimephales promelas	72	-----
14	71	Bluegill, Lepomis macrochirus	56	-----
		Redear sunfish, Lepomis microlophus	83	-----
		Green sunfish, Lepomis cyanellus	76	-----
13	55	Channel catfish, Ictalurus punctatus	46	-----
		Black bullhead, Ictalurus melas	64	-----
12	40	Yellow perch, Perca flavescens	40	-----

Table I2. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
11	35	Brook trout, <i>Salvelinus fontinalis</i>	44	-----
		Lake trout, <i>Salvelinus namaycush</i>	28	-----
10	33	Rainbow trout, <i>Oncorhynchus mykiss</i>	26	-----
		Coho salmon, <i>Oncorhynchus kisutch</i>	36	-----
		Chinook salmon, <i>Oncorhynchus tsawytscha</i>	40	-----
9	32	Largemouth bass, <i>Micropterus salmoides</i>	32	-----
8	26.11	Amphipod, <i>Gammarus fasciatus</i>	10.49	-----
		Amphipod, <i>Gammarus lacustris</i>	65	-----
7	20	Damselfly, <i>Lestes congener</i>	20	-----
6	20	Crawling water beetle, <i>Peltodytes sp.</i>	20	-----
5	13	Brown trout, <i>Salmo trutta</i>	13	-----
4	10	Isopod, <i>Asellus brevicaudus</i>	10	-----
3	3.3	Snail, <i>Lymnaea stagnalis</i>	3.3	-----
2	3	Backswimmer, <i>Notonecta undulata</i>	3	-----
1	2.1	Stonefly, <i>Pteronarcys californicus</i>	2.1	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

FAV = 1.903 ug/L

CMC = FAV/2 = 0.9515 ug/L

References

- Bluzat, R., and J. Senge. 1979. Effets de Trois insecticides (lindane, Fenthion and carbaryl): Toxicite' aigue sur gnatre espece d'invertebres limneques: Toxicite' chronique chez le mollusque pulmone' Lymnea. Environ. Pollut. 18:51-70.
- Federle, P.F., and W.J. Collins. 1976. Insecticide Toxicity to Three Insects from Ohio Ponds. Ohio J. Sci. 76:19-24.
- Hermens, J., H. Canton, M. Stezger, and R. Wagman. 1984. Joint Effects of a Mixture of 14 Chemicals on Mortality and Inhibition of Reproduction of Daphnia magna. Aquatic Toxicol. 5:315-322.
- Macek, K.J., and W.A. McAllister. 1970. Insecticide Susceptibility of Some Common Fish Family Representatives. Trans. Am. Fish. Soc. 99:20-27.
- Mayer, F.L., and M.R. Ellersieck. 1986. Manual of Acute Toxicity: Interpretation and Database for 410 Chemicals and 66 Species of Freshwater Animals. U.S. Dept. Interior, Fish and Wildlife Service. Resource Pub. 160. Columbia, MO.
- Randall, W.F., W.H. Dennis, and M.C. Warner. 1979. Acute Toxicity of Dechlorinated DDT, Chlordane and Lindane to Bluegill (Lepomis macrochirus) and Daphnia magna. Bull. Environ. Contam. Toxicol. 31:459-466.
- Tooby, T.E., and F.J. Durbin. 1975. Lindane Residue Accumulation and Elimination in Rainbow Trout (Salmo gairdneri Richardson) and Roach (Rutilus rutilus Linnaeus). Environ. Pollut. 8:79-89.
- U.S. EPA. 1980. Ambient Water Quality Criteria for Hexachlorocyclohexane. EPA 440/5-80-054. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Mercury(II)

The new acceptable acute data for mercury(II) are given in Table J1; no new chronic data were used. These new data were used with those given in Tables 1 and 2 of the criteria document for mercury(II) (U.S. EPA 1985) to obtain the values given in Table J2.

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values given in Table J2, resulting in a FAV of 3.388 ug/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 1.694 ug/L as total recoverable mercury(II).

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). ACRs were given for two freshwater species and one saltwater species in U.S. EPA (1985). The ACR obtained with the more resistant fathead minnow was much higher than the other two. The ACR obtained with the saltwater mysid was 3.095 and was similar to the Species Mean Acute-Chronic Ratio of 4.498 for *Daphnia magna*. The FACR was calculated as the geometric mean of the two SMACRs and was 3.731. The $FCV = FAV/FACR = (3.388 \text{ ug/L}) / (3.731) = 0.9081 \text{ ug/L}$. This value did not need to be lowered to protect a commercially or recreationally important species. The CCC was 0.9081 ug/L as total recoverable mercury(II).

The SMACR of >649.2 for the fathead minnow (Table J2) was not used in the calculation of the FACR because this species is acutely resistant to mercury(II). This SMACR is the geometric mean of >646.2, which was based on a life-cycle test, and >652.2, which was based on an early life-stage test. These two ACRs are so large that the two chronic values of <0.26 and <0.23 ug/L are both lower than the CCC of 0.9081 ug/L. Because the high SMACR was based on two tests with a fish and the two low SMACRs were obtained with invertebrates, it is quite possible that other fishes have SMACRs close to 649.2. The following estimated

chronic values were obtained using Species Mean Acute Values from Table J2 and an estimated ACR of 649.2:

<u>Species</u>	<u>Species Mean Acute Value</u>	<u>Estimated Chronic Value</u>
Rainbow trout	275 ug/L	0.42 ug/L
Coho salmon	240 ug/L	0.37 ug/L
Bluegill	160 ug/L	0.25 ug/L

All three of these estimated chronic values are for important species and are more than a factor of two lower than the FCV of 0.9081 ug/L. In addition, the SMACR for the fathead minnow is greater than 649.2. Thus the CCC of 0.9081 ug/L might not adequately protect such important fishes as the rainbow trout, coho salmon, and bluegill.

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of mercury(II) does not exceed 0.9081 ug/L more than once every three years on the average and if the one-hour average concentration does not exceed 1.694 ug/L more than once every three years on the average. The concentration of 0.9081 ug/L might not adequately protect such important fishes as the rainbow trout, coho salmon, and bluegill.

Table J1. New Acute Values for Mercury(II)

Species	Method*	Acute Value (ug/L)	Reference
Cladoceran, Ceriodaphnia reticulata	S,U	2.9	Elnabarawy et al. 1986
Cladoceran, Daphnia magna,	S,U	9.6	Elnabarawy et al. 1986
Cladoceran, Daphnia pulex	S,U	3.8	Elnabarawy et al. 1986
Amphipod, Crangonyx pseudogracilis	S,U	1.0**	Martin and Holdich 1986
Midge, Chironomus riparius	S,M	750	Rossaro et al. 1986
Mosquitofish, Gambusia affinis	S,U	230	Paulose 1988
Walking catfish, Clarias batrachus	S,U	375	Kirubagaran and Joy 1988
Fathead minnow, Pimephales promelas	FT,M	172	Spehar and Fiandt 1986
Guppy, Poecilia reticulata	R,U	26	Khangarot and Ray 1987

* S = static, R = renewal, FT = flow-through, U = unmeasured, M = measured.

** Not used in the derivation of the criterion because the corresponding 48-hr LC50 is 470 ug/L, which is an unusually large decrease in the LC50 from 48 to 96 hours.

Table J2. Ranked Genus Mean Acute Values for Mercury(II)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
29	2000	Stonefly, <i>Acroneuria lycorias</i>	2000	-----
28	2000	Mayfly, <i>Ephemera subvaria</i>	2000	-----
27	2000	Caddisfly, <i>Hydropsyche betteni</i>	2000	-----
26	1200	Caddisfly, (Unidentified)	1200	-----
25	1200	Damselfly, (Unidentified)	1200	-----
24	1000	Worm, <i>Nais</i> sp.	1000	-----
23	1000	Mozambique tilapia <i>Tilapia mossambica</i>	1000	-----
22	406.2	Tubificid worm, <i>Spirosperma ferox</i>	330	-----
		Tubificid worm, <i>Spirosperma rikolskyi</i>	500	-----
21	375	Walking catfish, <i>Clarias batrachus</i>	375	-----
20	370	Snail, <i>Aplexa hypnorum</i>	370	-----
19	257	Coho salmon, <i>Oncorhynchus kisutch</i>	240	-----
		Rainbow trout, <i>Oncorhynchus mykiss</i>	275	-----
18	250	Tubificid worm, <i>Quistadrilus multisetosus</i>	250	-----
17	240	Tubificid worm, <i>Rhyacodrilus montana</i>	240	-----
16	203	Mosquitofish, <i>Gambusia affinis</i>	203	-----
15	180	Tubificid worm, <i>Limnodrilus hoffmeisteri</i>	180	-----

Table J2. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
14	163	Fathead minnow, <i>Pimephales promelas</i>	163	> 649.2**
13	160	Bluegill, <i>Lepomis macrochirus</i>	160	-----
12	140	Tubificid worm, <i>Tubifex tubifex</i>	140	-----
11	140	Tubificid worm, <i>Stylodrilus heringianus</i>	140	-----
10	122***	Midge, <i>Chironomus</i> sp.	20	-----
		Midge, <i>Chironomus riparius</i>	750	-----
9	100	Tubificid worm, <i>Varichaeta pacifica</i>	100	-----
8	80	Tubificid worm, <i>Branchiura sowerbyi</i>	80	-----
7	80	Snail, <i>Amnicola</i> sp.	80	-----
6	50	Crayfish, <i>Orconectes limosus</i>	50	-----
5	28	Guppy, <i>Poecilia reticulata</i>	28	-----
4	20	Crayfish, <i>Faxonella clypeatus</i>	20	-----
3	10	Amphipod, <i>Gammarus</i> sp.	10	-----
2	3.3	Cladoceran, <i>Daphnia magna</i>	3.7	4.498
		Cladoceran, <i>Daphnia pulex</i>	2.9	-----
1	2.9	Cladoceran, <i>Ceriodaphnia reticulata</i>	2.9	-----

- * Ranked from most resistant to most sensitive based on Genus Mean Acute Value.
- ** Not used in the calculation of the Final Acute-Chronic Ratio.
- *** This GMAV was not set equal to the lowest SMAV because the species was not identified and so might have been *C. riparius*.

$$\text{FAV} = 3.388 \text{ ug/L}$$

$$\text{CMC} = \text{FAV}/2 = 1.694 \text{ ug/L}$$

$$\text{FACR} = 3.731$$

$$\text{FCV} = \text{FAV}/\text{FACR} = (3.388 \text{ ug/L})/(3.731) = 0.9081 \text{ ug/L} = \text{CCC}$$

The CCC of 0.9081 ug/L might not adequately protect such important fishes as the rainbow trout, coho salmon, and bluegill (see above).

References

- Elnabarawy, M.T., A.N. Welter, and R.R. Robideau. 1986. Relative Sensitivity of Three Daphnid Species to Selected Organic and Inorganic Chemicals. *Environ. Toxicol. Chem.* 5:393-398.
- Khengarot, B.S., and P.K. Ray. 1987. Studies on the Acute Toxicity of Copper and Mercury Alone and in Combination to the Common Guppy, *Poecilia reticulata* (Peters). *Arch. Hydrobiol.* 110:303-314.
- Kirubakaran, R., and K.P. Joy. 1988. Toxic Effects of Three Mercurial Compounds on Survival and Histology of the Kidney of the Catfish *Clarias batrachus* (L.). *Ecotoxicol. Environ. Safety* 15:171-179.
- Martin, T.R., and D.M. Holdich. 1986. The Acute Lethality of Heavy Metals to Peracarid Crustaceans (with Particular Reference to Fresh-Water Asellids and Gammarids). *Water Res.* 20:1137-1147.
- Paulose, P.V. 1988. Comparative Study of Inorganic and Organic Mercury Poisoning on Selected Freshwater Organisms. *J. Environ. Biol.* 9:203-206.
- Rossaro, B., G.F. Gaggino, and R. Marchetti. 1986. Accumulation of Mercury in Larvae and Adults, *Chironomus riparius* (Meigen). *Bull. Environ. Contam. Toxicol.* 37:402-406.
- Spehar, R.L., and J.T. Flandt. 1986. Acute and Chronic Effects of Water Quality Criteria-Based Metal Mixtures on Three Aquatic Species. *Environ. Toxicol. Chem.* 5:917-931.
- U.S. EPA. 1985. Ambient Water Quality Criteria for Mercury - 1984. EPA 440/5-84-026. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Nickel

The new acceptable acute data for nickel are given in Table K1; no new acceptable chronic data were found. These data were used with those given in Tables 1 and 2 of the criteria document for nickel (U.S. EPA 1986) to obtain the values given in Table K2. Some of the SMAVs in Table K2 differ from those given in Table 3 in U.S. EPA (1986) because preference was given to "FT,M" tests in Table K2. Because the toxicity of nickel is hardness-dependent, all acute values in Table K2 have been adjusted to a hardness of 50 mg/L.

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values in Table K2, resulting in an FAV of 522 ug/L at a hardness of 50 mg/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 261 ug/L, as total recoverable nickel, at a hardness of 50 mg/L. The CMC was related to hardness using the slope of 0.846 that was derived in U.S. EPA (1986):

$$\text{CMC} = e^{0.846 (\ln \text{hardness}) + 2.255}$$

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). SMACRs were available for two freshwater species and one saltwater species (U.S. EPA 1986). The saltwater ACR was 5.478 and the three are within a factor of 6.5. The FACR was calculated as the geometric mean of the three ACRs and was 17.99. The FCV = FAV/FACR = (522 ug/L)/(17.99) = 29.02 ug/L at a hardness of 50 mg/L. This value did not need to be lowered to protect a commercially or recreationally important species. Thus the CCC was 29.02 ug/L, as total recoverable nickel, at a hardness of 50 mg/L. The CCC was related to hardness using the slope of 0.846:

$$\text{CCC} = e^{0.846 (\ln \text{hardness}) + 0.0584}$$

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of nickel does not exceed the numerical value (in ug/L) given by the equation

$$CCC = e^{0.846(\ln \text{hardness}) + 0.0584}$$

more than once every three years on the average and if the one-hour average concentration does not exceed the numerical value (in ug/L) given by the equation

$$CMC = e^{0.846(\ln \text{hardness}) + 2.255}$$

more than once every three years on the average.

Table K1. New Acute Values for Nickel

Species	Method*	Hardness (mg/L as CaCO ₃)	Acute Value (ug/L)	Adjusted Acute Value (ug/L)**	Reference
Snail, <i>Physa gyrina</i>	FT,U	26	239	416	Nebeker et al. 1986
Amphipod, <i>Crangonyx pseudogracilis</i>	S,U	50	66,100	66,100	Martin and Holdich 1986
Midge, (1st instar) <i>Chironomus riparis</i>	S,U	55	72,400	66,791	Powlesland and George 1986
Midge, (1st instar) <i>Chironomus riparis</i>	S,U	55	81,300	75,002	Powlesland and George 1986
Midge, (1st instar) <i>Chironomus riparis</i>	S,U	55	84,900	78,323	Powlesland and George 1986
Midge, (2nd instar) <i>Chironomus riparis</i> ***	S,U	55	184,000	169,746	Powlesland and George 1986
Midge, (2nd instar) <i>Chironomus riparis</i> ***	S,U	55	150,000	138,380	Powlesland and George 1986
Midge, (2nd instar) <i>Chironomus riparis</i> ***	S,U	55	174,000	160,521	Powlesland and George 1986

* S = static, FT = flow-through, U = unmeasured.

** Adjusted to a hardness of 50 mg/L using a slope of 0.846.

*** Not used in the calculation of the SMAV because data were available for a more sensitive life stage.

Table K2. Ranked Genus Mean Acute Values for Nickel

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
21	73208	Midge, <i>Chironomus riparis</i>	73208	-----
20	66100	Amphipod, <i>Crangonyx pseudogracilis</i>	66100	-----
19	43250	Banded killifish, <i>Fundulus diaphanus</i>	43250	-----
18	40460	Stonefly, <i>Acroneuria lycorias</i>	40460	-----
17	30200	Caddisfly Unidentified sp.	30200	-----
16	21320	Goldfish, <i>Carassius auratus</i>	21320	-----
15	21200	Damselfly, Unidentified sp.	21200	-----
14	14100	Worm, <i>Nais</i> sp.	14100	-----
13	13380	Rainbow trout, <i>Oncorhynchus mykiss</i>	13380	-----
12	13000	Amphipod, <i>Gammarus</i> sp.	13000	-----
11	12770	Snail, <i>Amnicola</i> sp.	12770	-----
10	12756	Pumpkinseed, <i>Lepomis gibbosus</i>	7544	-----
		Bluegill, <i>Lepomis macrochirus</i>	21570	-----
9	12180	American eel, <i>Anguilla rostrata</i>	12180	-----
8	9839	Common carp, <i>Cyprinus carpio</i>	9839	-----
7	9661	Guppy, <i>Poecilia reticulata</i>	9661	-----

Table K2. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
6	8697	White perch, <i>Morone americana</i>	12790	-----
		Striped bass, <i>Morone saxatilis</i>	5914	-----
5	6707	Fathead minnow, <i>Pimephales promelas</i>	6707	35.58
4	4636	Mayfly, <i>Ephemerella subvaria</i>	4636	-----
3	4312	Rock bass, <i>Ambloplites rupestris</i>	4312	-----
2	1500	Cladoceran, <i>Daphnia pulicaria</i>	2042	-----
		Cladoceran, <i>Daphnia magna</i>	1102	29.86
1	416	Snail, <i>Physa gyrina</i>	416	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

** At hardness = 50 mg/L.

At hardness = 50 mg/L:

$$FAV = 522 \text{ ug/L}$$

$$CMC = FAV/2 = 261 \text{ ug/L}$$

As a function of hardness:

$$CMC = e^{0.846(\ln \text{ hardness}) + 2.255}$$

$$FACR = 17.99$$

At hardness = 50 mg/L:

$$FCV = FAV/FACR = (522 \text{ ug/L}) / (17.99) = 29.02 \text{ ug/L} = CCC$$

As a function of hardness:

$$CCC = e^{0.846(\ln \text{ hardness}) + 0.0584}$$

References

Martin, T.R., and D.M. Holdich. 1986. The Acute Lethal Toxicity of Heavy Metals to Peracarid Crustaceans (with Particular Reference to Fresh-water Asellids and Gammarids). *Water Res.* 20:1137-1147.

Nebeker, A.V., A. Stinchfield, C. Savonen, and G.A. Chapman. 1986. Effects of Copper, Nickel, and Zinc on Three Species of Oregon Freshwater Snails. *Environ. Toxicol. Chem.* 5:807-811

Powlesland, C., and J. George. 1986. Acute and Chronic Toxicity of Nickel to Larvae of *Chironomus riparis* (Meigan). *Environ. Poll., (Series A)*. 42:47-64.

U.S. EPA. 1987. Ambient Aquatic Life Water Quality Criteria for Nickel. EPA 440/5-86-004. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Parathion

No new acceptable acute or chronic data for parathion were found. Therefore, the data given in Tables 1 and 2 of the criteria document for parathion (U.S. EPA 1985) were used to obtain the values given in Table L1.

Criterion Maximum Concentration (CMC)

Some of the Genus Mean Acute Values given in Table 3 of U.S. EPA (1985) were changed because of the new taxonomy for salmonids and because only one value was calculated for the genus Chironomus; these changes did not affect the FAV. The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values given in Table L1, resulting in a FAV of 0.1299 ug/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 0.06495 ug/L.

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). Three Species Mean ACRs were available (Table L1). The ACRs obtained with the resistant fishes were much higher than that obtained with the sensitive cladoceran. To make the FACR appropriate for sensitive species, it was set equal to the ACR of 10.10 obtained with the cladoceran. The $FCV = FAV/FACR = (0.1299 \text{ ug/L})/(10.10) = 0.01286 \text{ ug/L}$. This value did not need to be lowered to protect a commercially or recreationally important species. The CCC was 0.01286 ug/L.

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of parathion does not exceed 0.01286 ug/L more than once every three years on the average and if the one-hour average concentration does not exceed 0.06495 ug/L more than once every three years on the average.

Table L1. Ranked Genus Mean Acute Values for Parathion

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
31	5,230	Tubificid worm, Tubifex sp.	5,230	-----
30	5,230	Tubificid worm, Limnodrilus sp.	5,230	-----
29	2,650	Channel catfish, Ictalurus punctatus	2,650	-----
28	2,223	Goldfish, Carassius auratus	2,223	-----
27	1,838	Brook trout, Salvelinus fontinalis	1,760	-----
		Lake trout Salvelinus namaycush	1,920	-----
26	1,510	Brown trout, Salmo trutta	1,510	-----
25	1,486	Cutthroat trout, Oncorhynchus clarki	1,560	-----
		Rainbow trout, Oncorhynchus gairdneri	1,415	-----
24	1,130	Isopod, Asellus brevicaudus	1,130	-----
23	1,000	Western chorus frog, Pseudacris triseriata	1,000	-----
22	839.6	Fathead minnow, Pimephales promelas	839.6	79.45**
21	688.7	Green sunfish, Lepomis cyanellus	930	-----
		Bluegill, Lepomis macrochirus	510	2121**
20	620	Largemouth bass, Micropterus salmoides	620	-----
19	320	Mosquitofish, Gambusia affinis	320	-----
18	<250	Crayfish, Procambarus sp.	<250	-----

Table L1. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
17	56	Guppy, <i>Poecilia reticulata</i>	56	-----
16	15	Mayfly, <i>Hexagenia bilineata</i>	15	-----
15	7.0	Beetle, <i>Peltodytes</i> spp.	7.0	-----
14	5.4	Stonefly, <i>Pteronarcys californica</i>	5.4	-----
13	4.2	Stonefly, <i>Pteronarcella badia</i>	4.2	-----
12	3.0	Damselfly, <i>Lestes congener</i>	3.0	-----
11	2.9	Stonefly, <i>Acroneuria pacifica</i>	2.9	-----
10	2.739	Prawn, <i>Palaemonetes kadiakensis</i>	2.739	-----
9	2.227	Mayfly, <i>Cloeon dipterum</i>	2.227	-----
8	1.697***	Midge, <i>Chironomus tentans</i>	31	-----
		Midge, <i>Chironomus riparius</i>	1.697	-----
7	1.5	Stonefly, <i>Claassenia sabulosa</i>	1.5	-----
6	1.127	Amphipod, <i>Gammarus fasciatus</i>	0.3628	-----
		Amphipod, <i>Gammarus lacustris</i>	3.5	-----
5	0.8944	Phantom midge, <i>Chaoborus</i> sp.	0.8944	-----

Table L1. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
4	0.7746	Cladoceran, Daphnia magna	1.0	10.10
		Cladoceran, Daphnia pulex	0.60	-----
3	0.64	Damselfly, Ischnura verticalis	0.64	-----
2	0.47	Cladoceran, Simocephalus serrulatus	0.47	-----
1	0.04	Crayfish, Orconectes nais	0.04	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

** Not used in the calculation of the Final Acute-Chronic Ratio.

*** This GMAV was set equal to the lower SMAV due to the large range in the SMAVs in this genus.

$$FAV = 0.1299 \text{ ug/L}$$

$$CMC = FAV/2 = 0.06495 \text{ ug/L}$$

$$FACR = 10.10$$

$$FCV = FAV/FACR = (0.1299 \text{ ug/L}) / (10.10) = 0.01286 \text{ ug/L} = CCC$$

References

U.S. EPA.: 1986. Ambient Aquatic Life Water Quality Criteria for Parathion - 1986. EPA 440/5-86-007. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Pentachlorophenol

No new acceptable acute or chronic data for pentachlorophenol were found. Therefore, the data given in Tables 1 and 2 of the criteria document for pentachlorophenol (U.S. EPA 1986) were used to obtain the values given in Table M1. Because the toxicity of pentachlorophenol is pH-dependent, all acute values in Table M1 have been adjusted to a pH of 6.5.

Criterion Maximum Concentration (CMC)

Some of the Genus Mean Acute Values given in Table 3 of U.S. EPA (1985) were changed because of the new taxonomy for salmonids and because the values for *Jordanella floridae* and *Rana catesbeiana* had been incorrectly adjusted to a pH of 6.5 and because the SMAV for *Gammarus pseudolimnaeus* had been calculated incorrectly. The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values given in Table M1, resulting in a FAV of 10.56 ug/L at a pH of 6.5. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 5.28 ug/L at a pH of 6.5. The CMC was related to pH using the slope of 1.005 that was derived in U.S. EPA (1986):

$$CMC = e^{1.005(\text{pH}) - 4.869}$$

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). Six Species Mean ACRs were available (Table M1), but two of them were "greater than" values. The range of the other four was less than a factor of 6. The FACR was calculated as the geometric mean of the four similar SMACRs and was 2.608. The FCV = FAV/FACR = (10.56 ug/L)/(2.608) = 4.049 ug/L at a pH of 6.5. This value did not need to be lowered to protect a commercially or recreationally important species. The CCC was 4.049 ug/L at a pH of 6.5. The CCC was related to pH using the slope of 1.005:

$$CCC = e^{1.005(\text{pH}) - 5.134}$$

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of pentachlorophenol does not exceed the numerical value (in ug/L) given by the equation

$$CCC = e^{1.005(\text{pH}) - 5.134}$$

more than once every three years on the average and if the one-hour average concentration does not exceed the numerical value (in ug/L) given by the equation

$$CMC = e^{1.005(\text{pH}) - 4.869}$$

more than once every three years on the average.

Table M1. Ranked Genus Mean Acute Values for Pentachlorophenol

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
32	>43920	Crayfish, <i>Orconectes immunis</i>	>43920	-----
31	11260	Midge, <i>Tanytarsus dissimilis</i>	11260	-----
30	10610	Sciomyzid, <i>Sepedon fuscipennis</i>	10610	-----
29	417.7	Tubificid worm, <i>Rhyacodrilus montana</i>	417.7	-----
28	408.2	Tubificid worm, <i>Stylodrilus heringianus</i>	408.2	-----
27	403.2	Snail, <i>Gilila altilis</i>	403.2	-----
26	361.6	Tubificid worm, <i>Spirosperma ferox</i>	239.5	-----
		Tubificid worm, <i>Spirosperma nikoiskyl</i>	545.8	-----
25	317.5	Tubificid worm, <i>Quistadrilus multisetosus</i>	317.5	-----
24	306.7	Flagfish, <i>Jordanella floridae</i>	306.7	-----
23	224.2	Tubificid worm, <i>Tubifex tubifex</i>	224.2	-----
22	195.4	Guppy, <i>Poecilia reticulata</i>	195.4	-----
21	182.5	Tubificid worm, <i>Limnodrilus hoffmeisteri</i>	182.5	-----
20	172.1	Amphipod, <i>Crangonyx pseudogracilis</i>	172.1	-----
19	155.9	Tubificid worm, <i>Branchiura sowerbyi</i>	155.9	-----
18	132.1	Snail, <i>Physa gyrina</i>	132.1	>10.27***

Table M1. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
17	105.0	Largemouth bass, <i>Micropterus salmoides</i>	105.0	-----
16	91.48	Amphipod, <i>Gammarus pseudolimnaeus</i>	91.48	-----
15	87.48	Amphipod, <i>Hyaella azteca</i>	87.48	-----
14	78.10	Cladoceran, <i>Daphnia pulex</i>	90.83	-----
		Cladoceran, <i>Daphnia magna</i>	67.15	2.5
13	67.13	Cladoceran, <i>Ceriodaphnia reticulata</i>	67.13	>15.79***
12	65.53	Goldfish, <i>Carassius auratus</i>	65.53	-----
11	63.11	Fathead minnow, <i>Pimephales promelas</i>	63.11	4.535
10	60.50	Mosquitofish, <i>Gambusia affinis</i>	60.50	-----
9	60.43	Snail, <i>Aplexa hypnorum</i>	60.43	-----
8	58.47	Tubificid worm, <i>Varichaeta pacifica</i>	58.47	-----
7	57.72	Cladoceran, <i>Simocephalus vetulus</i>	57.72	0.8945
6	56.41	Bluegill, <i>Lepomis macrochirus</i>	56.41	-----
5	34.13	Brook trout, <i>Salvelinus fontinalis</i>	34.13	-----
4	33.91	Bullfrog, <i>Rana catesbeiana</i>	33.91	-----

Table M1. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
3	31.26	Rainbow trout, <i>Oncorhynchus mykiss</i>	35.34	4.564
		Coho salmon, <i>Oncorhynchus kisutch</i>	31.82	-----
		Sockeye salmon, <i>Oncorhynchus nerka</i>	32.85	-----
		Chinook salmon, <i>Oncorhynchus tshawytscha</i>	25.85	-----
2	26.54	Channel catfish, <i>Ictalurus punctatus</i>	26.54	-----
1	4.355	Common carp, <i>Cyprinus carpio</i>	4.355	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

** At pH = 6.5.

*** Not used in the calculation of the Final Acute-Chronic Ratio.

At pH = 6.5:

$$FAV = 10.56 \text{ ug/L}$$

$$CMC = FAV/2 = 5.28 \text{ ug/L}$$

As a function of pH:

$$CMC = e^{1.005(pH) - 4.869}$$

$$FACR = 2.608$$

At pH = 6.5:

$$FCV = FAV/FACR = (10.56 \text{ ug/L}) / (2.608) = 4.049 \text{ ug/L} = CCC$$

As a function of pH:

$$CCC = e^{1.005(pH) - 5.134}$$

References

U.S. EPA.: 1986. Ambient Aquatic Life Water Quality Criteria for Pentachlorophenol. EPA 440/5-88-009. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Selenium

The new acceptable acute data for selenium are given in Table N1; no new acceptable chronic data were found. These new data were used with those given in Tables 1 and 2 of the criteria document for selenium (U.S. EPA 1987) to obtain the values given in Tables N2 and N3.

Selenium(IV):

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values given in Table N2, resulting in a FAV of 371.8 ug/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 185.9 ug/L.

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). Four Species Mean ACRs were available (Table N2), but the one determined with the acutely resistant species was higher than the other three; the three were within a factor of 2.4. The FACR was calculated as the geometric mean of the three and was 7.998. The $FCV = FAV/FACR = (371.8 \text{ ug/L})/(7.998) = 46.49 \text{ ug/L}$. As in U.S. EPA (1987), this value was lowered to 27.6 ug/L to protect the commercially and recreationally important rainbow trout. The CCC was 27.6 ug/L.

Selenium(VI):

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values given in Table N3, resulting in a FAV of 25.066 ug/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 12.533 ug/L.

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). Three Species Mean ACRs were available (Table N3), and they increased as the acute sensitivities of the species increased. To make the FACR appropriate for sensitive species, it was set equal to the SMACR of 2.651 for the sensitive *Daphnia magna*. The $FCV = FAV/FACR = (25.066 \text{ ug/L})/(2.651) = 9.455 \text{ ug/L}$. This value did not need to be lowered to protect a commercially or recreationally important species. The CCC was 9.455 ug/L.

Total selenium:

As discussed in U.S. EPA (1987), field studies conducted on Belews Lake in North Carolina suggested that selenium might be more toxic to certain species of freshwater fish than had been observed in laboratory chronic toxicity tests. Based upon these field studies and some laboratory studies, the CCC for total selenium was set at 5 ug/L. The Final Acute-Chronic Ratio for total selenium was calculated as the geometric mean of the six ACRs in Tables N2 and N3 that are between 2.5 and 16.5 and was 7.737. The FAV was calculated by multiplying the CCC by the FACR and was 38.68 ug/L. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 19.34 ug/L as total recoverable selenium.

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the four-day average concentration of selenium does not exceed 5 ug/L more than once every three years on the average and if the one-hour average concentration does not exceed 19.34 ug/L more than once every three years on the average.

Table N1. New Acute Values for Selenium

Species	Method*	Chemical	Acute Value (ug/L)	Reference
Cladoceran, Daphnia magna	S,U	Na-selenite [Selenium(IV)]	680	Johnston 1987
Cladoceran, Daphnia magna	S,U	Na-selenate [Selenium(VI)]	750	Johnston 1987

* S = static, U = unmeasured.

Table N2: Ranked Genus Mean Acute Values for Selenium(IV)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
22	203000	Leech, <i>Nepheleopsis obscure</i>	203000	-----
21	42500	Midge, <i>Tanytarsus dissimilis</i>	42500	-----
20	35000	Common carp, <i>Cyprinus carpio</i>	35000	-----
19	34910	Snail, <i>Aplexa hypnorum</i>	34910	-----
18	30176	White sucker, <i>Catostomus commersoni</i>	30176	-----
17	28500	Bluegill, <i>Lepomis macrochirus</i>	28500	-----
16	26100	Goldfish, <i>Carassius auratus</i>	26100	-----
15	25934	Midge, <i>Chironomus plumosus</i>	25934	-----
14	24100	Snail, <i>Physa</i> sp.	24100	-----
13	13600	Channel catfish, <i>Ictalurus punctatus</i>	13600	-----
12	12600	Mosquitofish, <i>Gambusia affinis</i>	12600	-----
11	11700	Yellow Perch, <i>Perca flavescens</i>	11700	-----
10	10490	Rainbow Trout, <i>Oncorhynchus mykiss</i>	10490	141.5**
9	10200	Brook trout, <i>Salvelinus fontinalis</i>	10200	-----
8	6500	Flagfish, <i>Jordanella floridae</i>	6500	-----
7	2704	Amphipod, <i>Gammarus pseudolimnaeus</i>	2704	-----

Table N2. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
6	1796	Cladoceran, Daphnia magna	834	13.31
		Cladoceran, Daphnia pulex	3870	5.586
5	1783	Striped bass, Morone saxatilis	1783	-----
4	1700	Hydra, Hydra sp.	1700	-----
3	1601	Fathead minnow, Pimephales promelas	1601	6.881
2	<603.6	Cladoceran, Ceriodaphnia affinis	<603.6	-----
1	340	Amphipod, Hyaella azteca	340	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

** Not used in the calculation of the Final Acute-Chronic Ratio.

$$FAV = 371.8 \text{ ug/L}$$

$$CMC = FAV/2 = 185.9 \text{ ug/L}$$

$$FACR = 7.998$$

$$FCV = FAV/FACR = (371.8 \text{ ug/L}) / (7.998) = 46.49 \text{ ug/L}$$

Lowered to protect rainbow trout:

$$FCV = 27.6 \text{ ug/L} = CCC$$

Table N3. Ranked Genus Mean Acute Values for Selenium(VI)

Rank*	Genus Mean Acute Value (ug/L)	Species	Species Mean Acute Value (ug/L)	Species Mean Acute-Chronic Ratio
11	442000	Leech, Hephelopsis obscura	442000	-----
10	193000	Snail, Aplexa hypnorum	193000	-----
9	66000	Channel catfish, Ictalurus punctatus	66000	-----
8	63000	Bluegill, Lepomis macrochirus	63000	-----
7	47000	Rainbow trout, Oncorhynchus mykiss	47000	16.26
6	20000	Midge, Paratanytarsus parthenogeneticus	20000	-----
5	7300	Hydra, Hydra sp.	7300	-----
4	5500	Fathead minnow, Pimephales promelas	5500	9.726
3	760	Amphipod, Hyaella azteca	760	-----
2	550.1	Cladoceran, Daphnia magna	1230	2.651
		Cladoceran, Daphnia pulicaria	246	-----
1	65.38	Amphipod, Gammarus pseudolimnaeus	65.38	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

$$FAV = 25.066 \text{ ug/L}$$

$$CMC = FAV/2 = 12.533 \text{ ug/L}$$

$$FACR = 2.651$$

$$FCV = FAV/FACR = (25.066 \text{ ug/L}) / (2.651) = 9.455 \text{ ug/L} = CCC$$

References

Johnston, P.A. 1987. Acute Toxicity of Inorganic Selenium to *Daphnia magna* (Straus) and the Effect of Sub-acute Exposure upon Growth and Reproduction. *Aquatic Toxicol.* 10:335-352.

U.S. EPA. 1987. Ambient Aquatic Life Water Quality Criteria for Selenium. EPA 440/5-87-006. National Technical Information Service, Springfield, VA.

1995 UPDATE:
Freshwater Aquatic Life Criterion for Zinc

The new acceptable acute data for zinc are given in Table O1; no new acceptable chronic data were found. These data were used with those given in Tables 1 and 2 of the criteria document for zinc (U.S. EPA 1987) to obtain the values given in Table O2. Because the toxicity of zinc is hardness-dependent, all acute values in Table O2 have been adjusted to a hardness of 50 mg/L.

Criterion Maximum Concentration (CMC)

The Final Acute Value (FAV) was calculated using the four lowest Genus Mean Acute Values in Table O2, resulting in an FAV of 133.2 ug/L at a hardness of 50 mg/L. This value did not need to be lowered to protect a commercially or recreationally important species. The CMC was calculated by dividing the FAV by 2, resulting in a CMC of 66.6 ug/L, as total recoverable zinc, at a hardness of 50 mg/L. The CMC was related to hardness using the slope of 0.8473 that was derived in U.S. EPA (1987):

$$CMC = e^{0.8473(\ln \text{ hardness}) + 0.884}$$

Criterion Continuous Concentration (CCC)

Insufficient chronic toxicity data were available to calculate a Final Chronic Value (FCV) using the eight-family procedure. Sufficient chronic data were available to calculate a FCV by dividing the FAV by the Final Acute-Chronic Ratio (FACR). SMACRs were available for seven species (Table O2), but three were for resistant species and one was a "less than" value. The other three were within a factor of 10.4. The FACR was calculated as the geometric mean of the three SMACRs and was 1.994. According to the methodology, the FACR cannot be less than 2. The FCV = FAV/FACR = (133.2 ug/L)/(2) = 66.6 ug/L at a hardness of 50 mg/L. This value did not need to be lowered to protect a commercially or recreationally important species. Thus the CCC was 66.6 ug/L, as total recoverable zinc, at a hardness of 50 mg/L and equals the CMC. The CCC was related to hardness using the slope of 0.8473:

$$CCC = e^{0.8473(\ln \text{ hardness}) + 0.884}$$

When it equals the CMC, the CCC is irrelevant because the CMC has a shorter averaging period.

The Criterion

The procedures described in the methodology indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms should not be affected unacceptably if the one-hour average concentration of zinc does not exceed the numerical value (in ug/L) given by the equation

$$\text{CMC} = e^{0.8473 (\ln \text{hardness}) + 0.884}$$

more than once every three years on the average.

Table O1. New Acute Values for Zinc

Species	Method*	Hardness (mg/L as CaCO ₃)	Acute Value (ug/L)	Adjusted Acute Value (ug/L)**	Reference
Frog, Xenopus laevis	S,M	100	34500	19176	Dawson et al. 1988
Cladoceran, Daphnia magna	S,U	300	1100	241	Berglind and Dave 1984

* S = Static, M = measured, U = unmeasured.

** Adjusted to a hardness of 50 mg/L using slope = 0.8473.

Table O2. Ranked Genus Mean Acute Values for Zinc.

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
36	88960	Damselfly, <i>Argia</i> sp.	88960	-----
35	19800	Amphipod, <i>Crangonyx pseudogracilis</i>	19800	-----
34	19176	Frog, <i>Xenopus laevis</i>	19176	-----
33	18400	Worm, <i>Nais</i> sp.	18400	-----
32	17940	Banded killifish, <i>Fundulus diaphanus</i>	17940	-----
31	16820	Snail, <i>Amnicola</i> sp.	16820	-----
30	13630	American eel, <i>Anguilla rostrata</i>	13630	-----
29	10560	Pumpkinseed, <i>Lepomis gibbosus</i>	18790	-----
		Bluegill, <i>Lepomis macrochirus</i>	5937	-----
28	10250	Goldfish, <i>Carassius auratus</i>	10250	-----
27	9712	Worm, <i>Lumbriculus variegatus</i>	9712	-----
26	8157	Isopod, <i>Asellus bicrenata</i>	5731	-----
		Isopod, <i>Asellus communis</i>	11610	-----
25	8100	Amphipod, <i>Gammarus</i> sp.	8100	-----
24	7233	Common carp, <i>Cyprinus carpio</i>	7233	-----
23	6580	Northern squawfish, <i>Ptychocheilus oregonensis</i>	6580	-----
22	6053	Guppy, <i>Poecilia reticulata</i>	6053	-----

Table O2. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
21	6000	Golden shiner, <i>Notemigonus crysoleucas</i>	6000	-----
20	5228	White sucker, <i>Catostomus commersoni</i>	5228	-----
19	4900	Asiatic clam, <i>Corbicula fluminea</i>	4900	-----
18	4341	Southern platyfish, <i>Xiphophorus maculatus</i>	4341	-----
17	3830	Fathead minnow, <i>Pimephales promelas</i>	3830	5.644***
16	3265	Isopod, <i>Lirceus alabamae</i>	3265	-----
15	2176	Atlantic salmon, <i>Salmo salar</i>	2176	-----
14	2100	Brook trout, <i>Salvelinus fontinalis</i>	2100	2.335***
13	1707	Bryozoan, <i>Lophopodella carteri</i>	1707	-----
12	1672	Flagfish, <i>Jordanella floridae</i>	1672	41.2***
11	1607	Bryozoan, <i>Plumatella emarginata</i>	1607	-----
10	1578	Snail, <i>Helisoma campanulatum</i>	1578	-----
9	1353	Snail, <i>Physa gyrina</i>	1683	-----
		Snail, <i>Physa heterostropha</i>	1088	-----
8	1307	Bryozoan, <i>Pectinatella magnifica</i>	1307	-----
7	>1264	Tubificid worm, <i>Limnodrilus hoffmeisteri</i>	>1264	-----

Table O2. (Cont.)

Rank*	Genus Mean Acute Value (ug/L)**	Species	Species Mean Acute Value (ug/L)**	Species Mean Acute-Chronic Ratio
6	931.3	Rainbow trout, Oncorhynchus mykiss	689.3	1.554
		Coho salmon, Oncorhynchus kisutch	1628	-----
		Sockeye salmon, Oncorhynchus nerka	1502	<6.074***
		Chinook salmon, Oncorhynchus tshawytscha	446.4	0.7027
5	790	Mozambique tilapia, Tilapia mossambica	790	-----
4	299.8	Cladoceran, Daphnia magna	355.5	7.26
		Cladoceran, Daphnia pulex	252.9	-----
3	227.8	Longfin dace, Agosia chrysogaster	227.8	-----
2	119.4	Striped bass, Morone saxatilis	119.4	-----
1	93.95	Cladoceran, Ceriodaphnia dubia	174.1	-----
		Cladoceran, Ceriodaphnia reticulata	50.70	-----

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value.

** At hardness = 50 mg/L.

*** Not used in the calculation of the Final Acute-Chronic Ratio.

At hardness = 50 mg/L:

FAV = 133.2 ug/L

CMC = FAV/2 = 66.6 ug/L

As a function of hardness:

$$\therefore \text{CMC} = e^{0.8473(\ln \text{hardness}) + 0.884}$$

FACR = 1.994 but was raised to 2

At hardness = 50 mg/L:

$$\text{FCV} = \text{FAV}/\text{FACR} = (133.2 \text{ ug/L})/(2) = 66.6 \text{ ug/L} = \text{CCC}$$

As a function of hardness:

$$\text{CCC} = e^{0.8473(\ln \text{hardness}) + 0.884}$$

References

Berglind, R., and G. Dave. 1984. Acute Toxicity of Chromate, DDT, PCP, TPBS, and Zinc to *Daphnia magna* Cultured in Hard and Soft Water. *Bull. Environ. Contam. Toxicol.* 33:63-68.

Dawson, D.A., E.F. Stebler, S.L. Burks, and J.A. Bantle. 1988. Evaluation of the Developmental Toxicity of Metal-Contaminated Sediments Using Short-Term Fathead Minnow and Frog Embryo-Larval Assays. *Environ. Toxicol. Chem.* 7:27-34.

U.S. EPA. 1987. Ambient Aquatic Life Water Quality Criteria for Zinc. EPA 440/5-87-003. National Technical Information Service, Springfield, VA.



